

**Indiana Lake Water Quality Assessment Report  
For 2015-2018**



**Prepared by:**

**Melissa Laney, Sarah Powers, Cory Sauvé  
School of Public & Environmental Affairs  
Indiana University  
Bloomington, Indiana**

**Prepared for:**

**Indiana Department of Environmental Management  
Office of Water Quality  
Indianapolis, Indiana**

**January 2019**

Cover: Crooked Lake, Whitley County. September 2015

## Acknowledgments

This report represents four years of traversing the State of Indiana, reading maps, searching for lake access, extracting stuck trailers, seemingly endless washing of laboratory glassware, and most importantly, having the opportunity to visit and sample the beautiful and varied lakes of Indiana. This work required the dedicated efforts of many people. Therefore, it is with gratitude that we recognize the superb efforts of the following SPEA graduate students who conducted the lake sampling and laboratory analyses of the water samples:

|                       |                   |                   |
|-----------------------|-------------------|-------------------|
| Heather Bearnese-Loza | Mitchell Latta    | Joao Palma Chacon |
| James Bennett         | Austin Linville   | Lauren Salvato    |
| Jake Berger           | Lori Lovell       | Cory Sauv         |
| Kristin Berger        | Thomas Miller     | Daniel Soebbing   |
| Karina Cardella       | Rowan Mitton      | Leigh Stevenson   |
| Gabby Ghreichi        | Courtney Mobilian | Cory Shumate      |
| Lucas Graham          | Kerry Neil        | Erica Walker      |
| Amy Hagerdon          | Bruna Oliveria    | Jason Wenning     |

This work was made possible by a grant from the U.S. EPA Section 319 Nonpoint Source Program administered by the Indiana Department of Environmental Management (IDEM). The IDEM Project Officers were Laura Crane, Chelsea Cottingham, and Jamie Hosier.

**Table of Contents**

INDIANA CLEAN LAKES PROGRAM ..... 9

    Lake Water Quality Assessment..... 9

    Water Quality Parameters Included in Lake Assessments ..... 11

        pH..... 11

        Conductivity..... 11

        Alkalinity ..... 12

        Phosphorus..... 12

        Nitrogen ..... 12

        Light Transmission ..... 13

        Dissolved Oxygen..... 13

        Secchi Disk Transparency..... 13

        Plankton ..... 14

        Chlorophyll-a ..... 14

LAKE CLASSIFICATION ..... 15

    Lake Origin Classification ..... 15

    Trophic Classification..... 16

    Trophic State Indices ..... 17

    The Carlson Trophic State Index ..... 17

    Ecoregion Descriptions..... 18

METHODS ..... 19

    Field Procedures..... 19

    Lab Procedures..... 20

RESULTS ..... 22

    Water Characteristics ..... 24

        pH, Conductivity, and Alkalinity..... 24

        SRP and TP..... 25

        NO<sub>3</sub>-N, NH<sub>3</sub>-N, and Org-N..... 26

        Chlorophyll-a and Phytoplankton..... 27

        Secchi Disk Transparency and Trophic State ..... 29

    Spatial Patterns..... 31

    Lake Type Characteristics..... 34

DISCUSSION ..... 40

    State of Indiana Lakes..... 40

|                          |    |
|--------------------------|----|
| Spatial Patterns.....    | 41 |
| Lake Type Patterns ..... | 41 |
| CONCLUSIONS.....         | 42 |
| REFERENCES .....         | 43 |
| APPENDICES .....         | 45 |

## List of Figures

Figure 1 – Cross-section of a lake experiencing summer thermal stratification. Adapted from Olem and Flock (1990).

Figure 2 – The pH scale compared with common solutions. Source: Addy et al. (2004).

Figure 3 – A simplified aquatic food chain.

Figure 4 – The Lake Michigan, Saginaw, and Erie lobes of the most recent glacial episode affecting northern Indiana.

Figure 5 – The Carlson Trophic State Index.

Figure 6 – Ecoregions of Indiana.

Figure 7 – Lakes assessed from 2015 to 2018.

Figure 8 – Surface area distribution for **(A)** all 329 lakes sampled from 2015 to 2018, and **(B)** the distribution of lakes under 500 hectares.

Figure 9 – Maximum depth distribution for 329 lakes sampled from 2015 to 2018.

Figure 10 – pH distribution for 329 lakes from 2015 to 2018 by **(A)** epilimnion and **(B)** hypolimnion.

Figure 11 – Conductivity and alkalinity distribution for 329 lakes sampled from 2015 to 2018 by **(A)** epilimnetic conductivity, **(B)** hypolimnetic conductivity, **(C)** epilimnetic alkalinity, and **(D)** hypolimnetic alkalinity.

Figure 12 – Soluble reactive phosphorus (SRP) distribution for 329 lakes sampled from 2015 to 2018 by **(A)** total distribution of epilimnetic SRP concentrations, **(B)** epilimnetic SRP concentrations under 0.05 mg/L, **(C)** all hypolimnetic SRP concentrations, and **(D)** hypolimnetic SRP concentrations under 0.60 mg/L.

Figure 13 – Total phosphorus (TP) distribution for 329 lakes sampled from 2015 to 2018 by **(A)** the total distribution of epilimnetic TP concentrations, **(B)** epilimnetic TP concentrations under 0.10 mg/L, **(C)** all hypolimnetic TP concentrations, and **(D)** hypolimnetic TP concentrations under 1.00 mg/L.

Figure 14 – Nitrate-nitrogen ( $\text{NO}_3\text{-N}$ ), ammonia-nitrogen ( $\text{NH}_3\text{-N}$ ), and organic-nitrogen (Org-N) distributions for 329 lakes sampled from 2015 to 2018 by **(A)** epilimnetic  $\text{NO}_3\text{-N}$ , **(B)** epilimnetic  $\text{NH}_3\text{-N}$ , **(C)** epilimnetic Org-N, **(D)** hypolimnetic  $\text{NO}_3\text{-N}$ , **(E)** hypolimnetic  $\text{NH}_3\text{-N}$ , and **(F)** hypolimnetic Org-N.

Figure 15 – Distribution of **(A)** mean plankton natural units, **(B)** mean plankton cells, **(C)** blue-green cell dominance, and **(D)** mean chlorophyll-a concentration by lake type for 329 lakes sampled from 2015 to 2018.

Figure 16 – Distribution of **(A)** mean plankton natural units, **(B)** mean plankton cells, **(C)** blue-green dominance, **(D)** mean chlorophyll-a concentrations by trophic state for 329 lakes sampled from 2015 to 2018.

Figure 17 – Secchi depth distribution for 329 lakes sampled from 2015 to 2018.

Figure 18 – Number of lakes sampled from 2015 to 2018 ( $n = 329$ ) by Carlson TSI [chl-a].

Figure 19 – Carlson TSI [TP] plotted against Carlson TSI [chl-a] for 329 lakes sampled from 2015 to 2018. The red line indicated the predicted relationship between TSI [TP] and TSI [chl-a].

Figure 20 – Location of lakes from 2015 to 2018 (n = 329) by Carlson TSI [chl-a] overlain on Indiana ecoregions (represented by color and corresponding number).

Figure 21 – Chlorophyll-a (chl-a) distribution by ecoregion for 329 lakes sampled from 2015 to 2018. Figure **(A)** illustrates the total distribution by a dot plot and **(B)** illustrates the same distribution with a box plot.

Figure 22 – Total phosphorus (TP) distribution by ecoregion for 320 lakes sampled from 2015 to 2018 by the **(A)** total TP distribution and the **(B)** TP distribution under 0.50 mg/L.

Figure 23 – Secchi depth distribution by ecoregion for 329 lakes sampled from 2015 to 2018 by the **(A)** total Secchi depth distribution and **(B)** Secchi depth values under 8 meters.

Figure 24 – Carlson TSI [chl-a] distribution by ecoregion for 329 lakes from 2015 to 2018. Figure **(A)** illustrates the total distribution by a dot plot and **(B)** illustrates the same distribution with a box plot. The dashed line indicates the TSI break between eutrophic and mesotrophic.

Figure 25 – Surface area distribution by lake type for 329 lakes sampled from 2015 to 2018 by the **(A)** total surface area distribution and **(B)** distribution under 850 hectares (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 26 – Maximum depth distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure **(A)** illustrates the total distribution by a dot plot and **(B)** illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 27 – Alkalinity distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total alkalinity distribution and **(B)** distribution under 350 mg CaCO<sub>3</sub>/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 28 – Conductivity distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure **(A)** illustrates the total distribution by a dot plot and **(B)** illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 29 – pH distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total pH distribution and **(B)** distribution from 6 to 9 pH units (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 30 – Nitrate-nitrogen (NO<sub>3</sub>-N) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total NO<sub>3</sub>-N distribution and **(B)** distribution under 0.80 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 31 – Ammonia-nitrogen (NH<sub>3</sub>-N) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total NH<sub>3</sub>-N distribution and **(B)** distribution under 2.50 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 32 – Total phosphorus (TP) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total TP distribution and **(B)** distribution under 0.40 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 33 – Chlorophyll-a (chl-a) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total chl-a distribution and **(B)** distribution under 80 ug/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 34 – Secchi depth distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total Secchi depth distribution and **(B)** distribution under 6 meters (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Figure 35 – TSI [chl-a] distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure **(A)** illustrates the total TSI [chl-a] distribution by a dot plot and **(B)** illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

## **INDIANA CLEAN LAKES PROGRAM**

The Indiana Clean Lakes Program was created in 1989 as a program within the Indiana Department of Environmental Management's (IDEM) Office of Water Management. The program is administered through a grant to Indiana University's School of Public and Environmental Affairs (SPEA) in Bloomington. The Indiana Clean Lakes Program is a comprehensive, statewide public lake management program having five components:

1. Public information and education
2. Technical assistance
3. Volunteer lake monitoring
4. Lake water quality assessment
5. Coordination with other state and federal lake programs.

This document is a summary of lake water quality assessment (LWQA) results for 2015 to 2018.

### **Lake Water Quality Assessment**

The goals of the LWQA include: (a) identifying water quality trends in individual lakes, (b) identifying lakes that need special management, and (c) tracking water quality improvements due to industrial discharge and runoff reduction programs (Jones 1996).

This program only samples public lakes that generally have boat trailer access from a public right-of-way. Public lakes are defined as those that have navigable inlets or outlets, or those that exist on or adjacent to public land. Sampling occurs in late June, July, and August of each year to coincide with the period of thermal stratification and the period of poorest annual water quality in lakes (Figure 1). Most Indiana lakes with maximum depths of 16 to 23 feet (5–7 m) or greater undergo thermal stratification during the summer. The warming of lake surface water by sunlight and higher air temperatures cause the water to become less dense. The less dense water will then rise above the cold, denser water at the lake's bottom. Summer wind and waves may not be strong enough to overcome the density differences between the surface and bottom waters and **thermal stratification** occurs. In a stratified lake, the surface waters (**epilimnion**) circulate and are well mixed throughout the summer while the bottom waters (**hypolimnion**) may stagnate because they are isolated from the surface. Thus, water characteristics in the epilimnion and hypolimnion of a given lake may be considerably different during stratification.




Figure 1 – Cross-section of a lake experiencing summer thermal stratification. Adapted from Olem and Flock (1990).

To account for potential differences between the epilimnion and hypolimnion of stratified lakes, water samples are collected from the top two meters of the surface and from one to two meters above the bottom. In addition, dissolved oxygen and temperature are measured at one-meter intervals from the surface to the bottom of each lake.

Lakes were randomized and selected from our list of all public lakes and impoundments having a) a minimum surface area of 5 acres, and b) a usable boat ramp. This process was similar to that used by the United States Environmental Protection Agency (USEPA) in the National Lakes Assessment (NLA) of 2007, 2012, and 2017. The resulting list contained a total of 329 lakes and impoundments. We randomize the candidate lake list each survey year. We sampled lakes from this list beginning with the first lake at the top and working downward until we had sampled 80 lakes each survey year, repeating the randomization for the next year. Using this sampling scheme, our 2015-2018 results should be statistically significant for the entire state and we could then better discuss lake water quality in Indiana.

The 329 lakes in our randomized pool are a small fraction of the 1475 lakes, reservoirs, and ponds in our master lake list for Indiana. However, many of these other lakes are private, smaller than 5 acres in surface area, and/or have no usable boat ramp. While the randomized sampling scheme allows us to gain a better understanding of Indiana lake quality each year, it is

possible that the sampling frequency for any given lake would create long gaps between individual lake surveys.

## Water Quality Parameters Included in Lake Assessments

Monitoring lakes requires many different parameters to be sampled. The parameters analyzed in this assessment include:

### *pH*

pH is the measure of the acidity of a solution of water. The pH scale commonly ranges from 0 to 14 (Figure 2). The scale is not linear but rather logarithmic. For example, a solution with a pH of 6.0 is ten times more acidic than a solution with a pH of 7.0. Pure water is said to be neutral, with a pH of 7.0. Water with a pH below 7.0 is considered acidic while water with pH greater than 7.0 is considered basic or alkaline. The pH of most natural waters in Indiana is between 6.5 and 8.0. However, acidic deposition may cause lower pH in susceptible waters and high phytoplankton productivity (which consumes CO<sub>2</sub>, a weak acid) can result in pH values exceeding 9.0.




Figure 2 – The pH scale compared with common solutions. Source: Addy et al. (2004).

### *Conductivity*

Conductivity is a numerical expression of an aqueous solution's capacity to carry an electric current. This ability depends on the presence of ions, their total concentration, mobility, valence, relative concentrations, and on the temperature of the liquid (APHA 2005). Solutions of most inorganic acids, bases, and salts are relatively good conductors. Conductivities of natural lakes in Indiana generally range from 50 to 1,000  $\mu\text{mhos/cm}$ , but the conductivity of old coal mine lakes can be as high as 3,000  $\mu\text{mhos/cm}$ . In contrast, the conductivity of distilled water is less than 1  $\mu\text{mhos/cm}$ . As conductivity is the inverse of resistance, the unit of conductance is the mho, or in low-conductivity natural waters, the micromho ( $\mu\text{mhos}$ ).

## *Alkalinity*

Alkalinity is the sum total of components in the water that tend to elevate the pH to the alkaline side of neutrality, and is expressed commonly as milligrams per liter as calcium carbonate (mg/L as CaCO<sub>3</sub>). Alkalinity is a measure of the *buffering capacity* (ability to resist changes in pH) of the water, and since pH has a direct effect on organisms as well as an indirect effect on the toxicity of certain pollutants in the water, the buffering capacity is important to water quality. Commonly occurring materials in water that increase alkalinity are carbonates, bicarbonates, phosphates, and hydroxides. Limestone bedrock and thick deposits of glacial till are good sources of carbonate buffering. Lakes within such areas are usually well-buffered.

## *Phosphorus*

Phosphorus is an essential plant nutrient and most often controls aquatic plant (algae and macrophyte) growth in freshwater. It is found in fertilizers, human and animal wastes, and yard waste. There is no atmospheric (vapor) form of phosphorus. Because there are few natural sources of phosphorus and the lack of an atmospheric cycle, phosphorus is often a *limiting nutrient* in aquatic systems. This means that the relative scarcity of phosphorus may limit the ultimate growth and production of algae and rooted aquatic plants. Therefore, management efforts often focus on reducing phosphorus input to a receiving waterway because: (a) it can be managed, and (b) reducing phosphorus can reduce algae production. Two common forms of phosphorus are:

***Soluble reactive phosphorus (SRP)*** – SRP is dissolved phosphorus readily usable by algae. SRP is often found in very low concentrations in phosphorus-limited systems where the phosphorus is tied up in the algae and cycled very rapidly. Sources of SRP include fertilizers, animal wastes, and septic systems.

***Total phosphorus (TP)*** – TP includes dissolved and particulate forms of phosphorus. TP concentrations greater than 0.03 mg/L (or 30g/L) can cause algal blooms in lakes and reservoirs.

## *Nitrogen*

Nitrogen is an essential plant nutrient found in fertilizers, human and animal wastes, yard waste, and the air. About 80 percent of the atmosphere is nitrogen gas. Nitrogen gas diffuses into water where it can be “fixed” (converted) by blue-green algae to ammonia for algal use. Nitrogen can also enter lakes and streams as inorganic nitrogen and ammonia. Because nitrogen can enter aquatic systems in many forms, there is an abundant supply of available nitrogen in these systems. The three common forms of nitrogen are:

**Nitrate ( $\text{NO}_3^-$ )** – Nitrate is an oxidized form of dissolved nitrogen that is converted to ammonia by algae under anoxic (low or no oxygen) conditions. It is found in streams and runoff when dissolved oxygen is present, usually in the surface waters.

**Ammonia ( $\text{NH}_4^+$ )** – Ammonia is a form of dissolved nitrogen that is readily used by algae. It is the reduced form of nitrogen and is found in water where dissolved oxygen is lacking such as in a eutrophic hypolimnion. Important sources of ammonia include fertilizers and animal manure. In addition, ammonia is produced as a by-product by bacteria as dead plant and animal matter are decomposed.

**Organic Nitrogen (Org-N)** – Organic nitrogen includes nitrogen found in plant and animal materials and may be in dissolved or particulate form. In the analytical procedures, total Kjeldahl nitrogen (TKN) was determined in 2015. Organic nitrogen is TKN minus ammonia. In 2016, the analytical procedures were changed, and total nitrogen (TN) was determined. Organic nitrogen is TN minus nitrate and ammonia.

### *Light Transmission*

This measurement uses a light meter (photocell) to determine the *rate* at which light transmission is diminished in the upper portion of the lake's water column. Another important light transmission measurement is the determination of the 1% light level. The 1% light level is the water depth at which one percent of surface light penetrates. The 1% light level is considered the lower limit of algal growth in lakes and this area and above is referred to as the *euphotic zone*.

### *Dissolved Oxygen*

Dissolved oxygen (DO) is the dissolved gaseous form of oxygen. It is essential for respiration of fish and other aquatic organisms. DO enters water by diffusion from the atmosphere and as a by-product of photosynthesis by algae and plants. The concentration of DO in epilimnetic waters continually equilibrates with the concentration of atmospheric oxygen to maintain 100 percent DO saturation. Excessive algae growth can over-saturate (greater than 100 percent saturation) the water with DO when the rate of photosynthesis is greater than the rate of oxygen diffusion to the atmosphere. Hypolimnetic DO concentration is typically low as there is no mechanism to replace oxygen that is consumed by respiration and decomposition. Fish need at least 3-5 mg/L of DO to survive.

### *Secchi Disk Transparency*

Secchi disk transparency refers to the depth to which a black and white Secchi disk can be seen in the lake water. Water clarity, as determined by a Secchi disk, is affected by two primary factors: algae and suspended particulate matter. Particulates (soil or dead leaves) may be introduced into the water by either runoff or sediments already on the bottom of the lake.

Erosion from construction sites, agricultural lands, and riverbanks all lead to increased sediment runoff. Bottom sediments can be resuspended by bottom-feeding fish such as carp, by motorboats, or by strong winds in shallow lakes.

### Plankton

Plankton are important members of the aquatic food web. The plankton include phytoplankton or algae (microscopic plants) and zooplankton (tiny shrimp-like animals that eat algae). The phytoplankton are primary producers that convert light energy from the sun to plant tissue through the process of photosynthesis. This forms the foundation of the aquatic food chain. Small microscopic shrimp-like crustaceans – called zooplankton – eat the phytoplankton. In turn, the zooplankton are extremely important food for young fish (Figure 3).

The phytoplankton are organized taxonomically largely by color. Important phyla (groups) include: Cyanobacteria (blue-green algae), Chlorophyta (green algae), Chrysophyta (yellow-brown algae), and Bacillariophyta (diatoms). The cyanobacteria are of particular interest to limnologists and lake users because members of this group are those that often form nuisance blooms and their dominance in lakes may indicate poor water conditions. Some species of cyanobacteria are known toxin producers.

### Chlorophyll-a

The plant pigments of algae consist of the chlorophylls (green color) and carotenoids (yellow color). Chlorophyll-a is the most dominant chlorophyll pigment in green algae (Chlorophyta), but is only one of several pigments in blue-green algae (Cyanophyta), yellow-brown algae (Chrysophyta), and others. Despite this, chlorophyll-a is often used as a direct estimate of algal biomass although it might underestimate the production of algae that contain multiple pigments.




Figure 3 – A simplified aquatic food chain.

## LAKE CLASSIFICATION

There are many factors that influence the condition of a lake including physical dimensions (*morphometry*), nutrient concentrations, oxygen availability, temperature, light, and fish species. In order to simplify the analysis of lakes, there are a variety of lake classifications that are used. Lake classifications serve to aid in the decision-making process, in prioritizing, and in creating public awareness. Lakes can be classified based on their origin, thermal stratification regime, or by trophic status.

### **Lake Origin Classification**

Hutchinson (1957) classified lakes according to how they were formed which resulted in 76 different classifications; the following are several important lake types in Indiana.

#### *Glacial Lakes*

As the glacier ice sheets moved south and then receded some 10,000 to 12,000 years ago, they created several types of lakes including scour lakes and kettle lakes. **Scour lakes** were formed when the sheet moved over the land creating a groove in the surface of the earth which later filled with meltwater. **Kettle lakes** were formed when large chunks of ice, deposited by the retreating glacier, left depressions in the thick deposits of *till* (sand and gravel ground up by the glacier) that covered the landscape. When the ice blocks melted the depressions filled in with water and lakes were formed. The majority of lakes in Indiana are kettle lakes including Lake Tippecanoe, the deepest lake (123 feet), and Lake Wawasee, the largest glacial lake (3,410 acres). Glacial lakes in Indiana are primarily in the north and are found between the western Valparaiso Morainal Area and the eastern Steuben Morainal Area where the Lake Michigan, Saginaw, and Erie lobes occurred (Figure 4). Glacial lakes are thus limited to this part of the state.

#### *Solution Lakes*

Solution lakes form when water collects in basins formed by the solution of limestone found in regions of karst topography. These lakes tend to be circular and are primarily found in the Mitchell Plain of southern Indiana.




Figure 4 – The Lake Michigan, Saginaw, and Erie lobes of the most recent glacial episode affecting northern Indiana.

### *Oxbow Lakes*

Oxbow lakes are formed from former river channels that have been isolated from the original river channel due to deposition of sedimentation or erosion. Oxbow lakes can be found throughout the state of Indiana.

### *Artificial Lakes*

Artificial lakes are created by humans due to excavation of a site or to damming a stream or river. Artificial lakes include ponds, strip pits, borrow pits, quarries, and reservoirs (Jones 1996). Reservoirs, also called impoundments, are typically elongated with many branches representing the tributaries of the former stream or river. Strip pits are coal mine lakes found in southwestern Indiana where coal mines are located. Many coal mine lakes formed when water filled the final cut excavated during surface mining. Borrow pits were originally excavated as a source of fill dirt for highway and other large construction projects. For our purposes, we aggregated strip pits, borrow pits, and quarry pits into a singular classification, surface mine lakes (SML).

### **Trophic Classification**

Trophic state is an indication of a lake's nutritional level or biological productivity. The following definitions are used to describe the trophic state of a lake:

*Oligotrophic* - lakes with clear waters, low nutrient levels (total phosphorus < 6 µg/L), supports few algae, hypolimnion has dissolved oxygen, and can support salmonids (trout and salmon).

*Mesotrophic* - water is less clear, moderate nutrient levels (total phosphorus 10-30 µg/L), support healthy populations of algae, less dissolved oxygen in the hypolimnion, and lack of salmonids.

*Eutrophic* - water transparency is less than 2 meters, high concentrations of nutrients (total phosphorus > 35 µg/L), abundant algae and weeds, lack of dissolved oxygen in the hypolimnion during the summer.

*Hypereutrophic* - water transparency less than 1 meter, extremely high concentrations of nutrients (total phosphorus > 80 µg/L), thick algal scum, dense weeds.

Eutrophication is the biological response observed in a lake caused by increased nutrients, organic material, and/or silt (Cooke et al. 1993). Nutrients enter the lake through runoff or through eroded soils to which they are attached. Increased nutrient concentrations stimulate the growth of aquatic plants. Sediments and plant remains accumulate at the bottom of the lake decreasing the mean depth of the lake. The filling-in of a lake is a natural process that usually occurs over thousands of years. However, this natural process can be accelerated by

human activities such as increased watershed erosion and increased nutrient loss from the land. Thus, **cultural eutrophication** can degrade a lake in as little as a few decades.

Although it is widely known that nutrients, especially phosphorus, are responsible for increased productivity, the concentration of nutrients alone cannot determine the trophic state of a lake. Other factors such as the presence of algae and weeds aid in the determination of the trophic status, and other factors such as light and temperature impact the growth of algae and weeds.

### Trophic State Indices

Due to the complex nature and variability of water quality data, a trophic state index (TSI) is used to aid in the evaluation of water quality data. A TSI assigns a numerical value to different levels of standard water quality measurements. The sum of these points for all parameters in the TSI represents the standardized trophic status of a lake that can be compared in different years or can be compared to other lakes. When using a TSI for comparison, it is important to not neglect the actual data as these data may help in explaining other differences between lakes. As with any index, when the data are reduced to a single number for a TSI, some information is lost.

### The Carlson Trophic State Index

The Carlson Trophic State Index, developed by Carlson (1977) is the most widely used TSI in the United States (Figure 5). Carlson used mathematical equations developed from the relationships observed between summer measurements of Secchi disk transparency, total phosphorus, and chlorophyll-a in north temperate lakes. With Carlson’s TSI, one parameter, Secchi disk transparency, total phosphorus, or chlorophyll-a, can be used to yield a TSI value for that lake. One parameter can also be used to predict the value of the other parameters. Values for the Carlson’s TSI range from 0 to 100 and each increase of 10 trophic points represents a doubling of algal biomass.

Not all lakes exhibit the same relationship between Secchi disk transparency, total phosphorus, and chlorophyll-a that Carlson’s lakes show. However, in these cases Carlson’s TSI gives valuable insight into the functioning of a particular lake.




Figure 5 – The Carlson Trophic State Index.

## Ecoregion Descriptions

The connection between lakes and their associated watersheds is evident in processes that include soil types, land slope, and watershed land use. These relationships can be expanded to a larger scale – the ecoregion – that incorporate these relationships across a larger geographic area. Omernik and Gallant (1988) defined ecoregions in the Midwest; the boundaries of these ecoregions were determined through the examination of land use, soils, and potential natural vegetation. These ecoregions have similar ecological properties throughout their range and these properties can influence lake water quality characteristics. Six ecoregions are present in Indiana (Figure 6). Descriptions of the ecoregions are as follows:

**Central Corn Belt Plains (#54):** This ecoregion covers 46,000 square miles of Indiana and Illinois. This ecoregion is primarily cultivated for feed crops, only 5 percent of the area is woodland. Crops and livestock are responsible for the nonpoint source pollution in this region.

**Eastern Corn Belt Plains (#55):** This ecoregion covers 31,800 square miles of Indiana, Ohio, and Michigan. Hardwood forests can thrive in this area; 75 percent of the land is used for crop production. Few natural lakes or reservoirs are located in this area.




Figure 6 – Ecoregions of Indiana.

**Southern Michigan/Northern Indiana Till Plain (#56):** This region covers 25,800 square miles of Michigan and Indiana. Oak-hickory forests are the dominant vegetation in this area; however, 25 percent of this area is urbanized.

**Huron/Erie Lake plain (#57):** This region covers 11,000 square miles of Indiana, Ohio, and Michigan. This area used to be occupied by forested wetlands; however, the primary use is now farming and 10 percent of this region is urbanized. There are no lakes in this region that could be assessed by the present study.

**Interior Plateau (#71):** This area occupies 56,000 square miles from Indiana and Ohio down to Alabama. Land is used for pasture, livestock, and crops. Woodlands and forests remain in this area. There are many quarries and coal mines in this area; however, there are few natural lakes.

**Interior River Lowland (#72):** This area covers 29,000 square miles in Indiana, Kentucky, Illinois, and Missouri. One third of this area is maintained as oak-hickory forest; other land uses include pasture, livestock, crops, timber, and coal mines. Water quality disturbances come from livestock, crops, and surface mining.

## **METHODS**

### **Field Procedures**

Water samples are collected from the epilimnion and hypolimnion, generally 1 meter below the surface and from 1-2 meters above the bottom of the lake. Epilimnetic water samples were collected using a 2-meter long integrated sampler that samples an undisturbed column of water from the surface to a depth of 2-meters. The sampler is emptied into a clean, rinsed pitcher where it is thoroughly mixed before filling the sample bottles. Water samples were taken for soluble reactive phosphorus (SRP), total phosphorus (TP), nitrate ( $\text{NO}_3^-$ ), ammonia ( $\text{NH}_4^+$ ), and total nitrogen (TN). SRP is filtered in the field using a 47  $\mu\text{m}$  membrane filter. Prior to sampling, the TP, nitrate/ammonia, and TN bottles are acidified with sulfuric acid ( $\text{H}_2\text{SO}_4$ ) resulting in a pH of the sample between 1 and 2.

Dissolved oxygen (DO), temperature, conductivity, and pH are measured using an In-Situ multi-parameter sonde. Measurements are taken at 1-meter intervals through the water column to the lake bottom.

Secchi disk transparency is measured by lowering a black and white disk through the water column until it is no longer visible. Light penetration is measured with a LiCor Spherical Quantum Sensor.

Phytoplankton were sampled using a 2-meter integrated sampler. The sampler is emptied into a clean, rinsed pitcher where it is thoroughly mixed before filling the sample bottles. The

phytoplankton samples were preserved with glutaraldehyde during post-sampling activities. Zooplankton were collected with a tow net through the whole water column, utilizing a 80-micron mesh on the net and bucket. Zooplankton samples were preserved with 95% ethyl alcohol.

Chlorophyll-a is collected with an integrated sampler that reaches to a 2-m depth. The apparatus is shut, retrieved, and poured into a pitcher. The sample is shaded and filtered with Whatman GF/F filter paper using a hand pump. The sample is filtered until the flow of water passing through the filter is minimal and the volume of sample filtered is then recorded. The filter paper is removed, placed in a bottle, and kept thoroughly chilled.

### **Lab Procedures**

SRP is determined using an ascorbic acid method and then measured colorimetrically on a spectrophotometer (APHA 2005). Prior to 2016, TP samples were analyzed using a nitric and sulfuric acid digestion to convert particulate phosphorus to dissolved phosphorus. After pH adjustment, the samples are analyzed for SRP.

$\text{NO}_3^-$  is analyzed using the cadmium reduction method (US EPA Method 353.3) using segmented flow analysis.  $\text{NH}_4^+$  is processed using in-line gas diffusion (US EPA Method 350.1) (US EPA 1993). TN samples are digested in an alkaline persulfate solution then processed as nitrite-nitrate using the cadmium reduction method. TP samples are digested in an alkaline persulfate solution releasing particulate bound phosphorus and analyzed using the ascorbid acid method. Segmented flow analysis is performed on an Alpkem Flow Solution Model 3570 autoanalyzer. TKN (2015 only) samples are first digested in hot acid before being analyzed with an autoanalyzer.

For zooplankton analysis, one milliliter of water is transferred to a Sedgwick-Rafter Cell for identification and enumeration. The entire cell is scanned and all zooplankton are counted. Whole water samples of phytoplankton were concentrated to insure sufficient cell density using Utermoehl settling chambers. Counts are made using a nanoplankton chamber (PhycoTech, Inc.) and a phase contrast light microscope (2015-2017). In 2018, whole water samples of phytoplankton were concentrated using vacuum filtration and HPMA mounting (PhycoTech 2018). Plankton identifications are made according to: Ward and Whipple (1959), Prescott (1982), Whitford and Schumacher (1984), Wehr and Sheatch (2003), and St. Amand (2010). After identification, the following parameters can be calculated:

1. **Natural Unit density (NU/L)** – this is the historic unit used for many years to quantify plankton in Indiana lakes. A natural unit represents a single organism, irregardless of whether the organism is single-celled or a multi-celled colonial form. The size range of natural units may be several orders of magnitude (100 – 1000x).
2. **Cell density (cells/mL)** – Counting and recording at the cell level is preferred by phycologists and limnologists today. Each phytoplankton cell can live and reproduce

independently of other cells, even in those taxa that aggregate in colonies. Public health warnings regarding toxigenic cyanobacteria are determined, in part, by cell densities.

3. **Blue-green dominance (%)** – This valuable variable is the percentage of a plankton population that is dominated by cyanobacteria. Since cyanobacteria are more likely to become a nuisance in aquatic systems, this simple indicator is still useful. Caution is necessary in interpreting this metric because dominance by cyanobacteria in a lake with a low density of phytoplankton does not necessarily indicate a problem in that lake.

For chlorophyll-a analysis, filters are placed in the freezer upon arriving to the lab. Once frozen, the filters are ground and chlorophyll-a is extracted using 90% aqueous acetone and measured using spectrophotometer. Samples are corrected for pheophyton pigments with dilute acid.

All sampling techniques and laboratory analytical methods were performed in accordance with procedures in APHA (2005). Details can be found in the Quality Assurance Protection Plan (QAPP).

## **RESULTS**

Information about the lakes sampled from 2015 to 2018 is included in Appendix A and B. Raw data for all lakes assessed are available on the Indiana Clean Lakes Program website at: <http://www.indiana.edu/~clp>.

### **Lakes Assessed**

We assessed a total of 329 lakes during this four-year period; 81 in 2015, 86 in 2016, 80 in 2017, and 82 in 2018 (Figure 7).




Figure 7 – Lakes assessed from 2015 to 2018.

Lake surface area ranged from 0.30 hectares (Skunk Lake) to 4353.75 hectares (Monroe Reservoir), with a median surface area of 26.31 ha (Figure 8). Twenty lakes had surface areas greater than 500 ha, while 64.7 percent (n = 213) of all lakes sampled were under 50 ha.




Figure 8 – Surface area distribution for **(A)** all 329 lakes sampled from 2015 to 2018, and **(B)** the distribution of lakes under 500 hectares.

Maximum depth ranged from 1.4 meters (Nasby Mill Pond) to 37 meters (Tippecanoe Lake), with a median of 9.1 meters (Figure 9). Natural lakes had the deepest median maximum depth (10.7 meters), followed by surface mine lakes (7.62 meters) and impoundments (6.4 meters).




Figure 9 – Maximum depth distribution for 329 lakes sampled from 2015 to 2018.

## Water Characteristics

### *pH, Conductivity, and Alkalinity*

Epilimnetic pH ranged from 5.8 to 9.4 for all lakes sampled. Schlamm Lake had the lowest epilimnetic pH of 5.8 and Bobcat Lake – a surface mine lake – had the highest epilimnetic pH of 9.4 (Figure 10). Median epilimnetic pH for all lakes was 7.9. Hypolimnetic pH was comparable to epilimnetic pH, with a median hypolimnetic pH of 7.2. Schlamm Lake and Bobcat Lake again represented the lowest and highest hypolimnetic pH, with a pH of 5.7 and 9.5, respectively.




Figure 10 – pH distribution for 329 lakes from 2015 to 2018 by (A) epilimnion and (B) hypolimnion.

Median epilimnetic and hypolimnetic values were comparable for both conductivity and alkalinity (Figure 11). Minimum epilimnetic and hypolimnetic conductivity values were 46 umhos/cm and 113 umhos/cm, respectively. Maximum hypolimnetic conductivity was also higher than maximum epilimnetic conductivity, with values of 467.4 and 2,800 umhos/cm, respectively. Both of these lakes are surface mine lakes – Trimble Lake and Hale Lake – located in Greene and Sullivan County. Median epilimnetic conductivity was 416.5 umhos/cm compared to 467.4 for hypolimnetic samples.

The median alkalinity concentration for epilimnetic samples was 143.5 mg CaCO<sub>3</sub>/L and 182 mg CaCO<sub>3</sub>/L for hypolimnetic samples. Sycamore Lake (343 mg CaCO<sub>3</sub>/L) and Airline Lake (182 mg CaCO<sub>3</sub>/L) represented the maximum values for both samples, and are both surface mine lakes located in Greene County.




Figure 11 – Conductivity and alkalinity distribution for 329 lakes sampled from 2015 to 2018 by (A) epilimnetic conductivity, (B) hypolimnetic conductivity, (C) epilimnetic alkalinity, and (D) hypolimnetic alkalinity.

*SRP and TP*

Epilimnetic SRP concentrations were generally low across all lakes, with a median concentration of 0.008 mg/L (Figure 12). Fifty-one lakes (15.5 percent) were at or below the method detection limit of 0.002 mg/L. However, Shakamak Lake in Sullivan County had a SRP concentration of 1.227 mg/L. Hypolimnetic SRP were higher than epilimnetic samples, with a median concentration of 0.160 mg/L. Only 16 lakes (4.9 percent) of lakes were at or below the method detection limit. Riddles Lake had the highest concentration of 1.489 mg/L.




Figure 12 – Soluble reactive phosphorus (SRP) distribution for 329 lakes sampled from 2015 to 2018 by (A) total distribution of epilimnetic SRP concentrations, (B) epilimnetic SRP concentrations under 0.05 mg/L, (C) all hypolimnetic SRP concentrations, and (D) hypolimnetic SRP concentrations under 0.60 mg/L.

Epilimnetic TP concentrations were lower compared to hypolimnetic samples, with median concentrations of 0.031 and 0.098 mg/L, respectively (Figure 13). Loomis Lake (Porter County) had the highest epilimnetic TP concentration of 0.609 mg/L and Airline Lake had the highest hypolimnetic concentration of 3.97 mg/L.




Figure 13 – Total phosphorus (TP) distribution for 329 lakes sampled from 2015 to 2018 by (A) the total distribution of epilimnetic TP concentrations, (B) epilimnetic TP concentrations under 0.10 mg/L, (C) all hypolimnetic TP concentrations, and (D) hypolimnetic TP concentrations under 1.00 mg/L.

### *NO<sub>3</sub>-N, NH<sub>3</sub>-N, and Org-N*

Nitrate-nitrogen median epilimnetic and hypolimnetic concentrations were similar, with concentrations of 0.014 and 0.013 mg/L, respectively (Figure 14). For the 329 lakes sampled, 77 lakes had epilimnetic nitrate-nitrogen concentrations below the method detection limit of 0.008 mg/L, and 74 lakes had hypolimnetic concentrations below the method detection limit.

Ammonia-nitrogen concentrations were higher in the hypolimnion than the epilimnion (Figure 14). The median epilimnetic ammonia-nitrogen concentration was 0.018 mg/L, with 21.3 percent of lakes sampled below the method detection limit of 0.014 mg/L. In contrast, the median hypolimnetic concentration was 0.748 mg/L with only 3 percent of lakes below the method detection limit. Airline Lake (Greene Co.) had the highest hypolimnetic concentration of 52.801 mg/L.

The median organic-nitrogen concentration was 0.260 mg/L (Figure 14). While North Twin Lake (LaGrange Co.) had the highest organic-nitrogen concentration of 8.542 mg/L, organic-nitrogen concentrations across the 329 lakes were more normally distributed compared to nitrate-nitrogen and ammonia-nitrogen.




Figure 14 – Nitrate-nitrogen (NO<sub>3</sub>-N), ammonia-nitrogen (NH<sub>3</sub>-N), and organic-nitrogen (Org-N) distributions for 329 lakes sampled from 2015 to 2018 by **(A)** epilimnetic NO<sub>3</sub>-N, **(B)** epilimnetic NH<sub>3</sub>-N, **(C)** epilimnetic Org-N, **(D)** hypolimnetic NO<sub>3</sub>-N, **(E)** hypolimnetic NH<sub>3</sub>-N, and **(F)** hypolimnetic Org-N.

### *Chlorophyll-a and Phytoplankton*

Chlorophyll-a concentrations ranged from 0.398 ug/L (Gambill Lake, Sullivan Co.) to 146.595 ug/L (Waveland Lake, Montgomery Co.), with a median concentration of 6.966 ug/L. Chlorophyll-a concentrations were highest in impoundments with a mean concentration of 17.90 ug/L, compared to 7.432 ug/L for natural lakes and 3.183 ug/L for surface mine lakes (Figure 15). Hypereutrophic lakes had the highest mean chlorophyll-a concentration of 59.269 ug/L, followed by eutrophic lakes (16.681 ug/L), mesotrophic lakes (4.298 ug/L), and oligotrophic lakes (1.263 ug/L) (Figure 16).




Figure 15 – Distribution of (A) mean plankton natural units, (B) mean plankton cells, (C) blue-green cell dominance, and (D) mean chlorophyll-a concentration by lake type for 329 lakes sampled from 2015 to 2018.




Figure 16 – Distribution of (A) mean plankton natural units, (B) mean plankton cells, (C) blue-green dominance, (D) mean chlorophyll-a concentrations by trophic state for 329 lakes sampled from 2015 to 2018.

Mean phytoplankton cell concentration was 71,954.74 cells/mL. Three lakes had concentrations exceeding 1 million cells/mL: Canada Lake (Porter Co.), Fish Lake (LaPorte Co.), and Sycamore Lake (Green Co.). Phytoplankton cell concentrations followed similar trends to that of chlorophyll-a in terms of lake type and trophic state. Mean cell concentrations were highest for impoundments (19,022 cells/mL), followed by natural lakes (13,480 cells/mL) and surface mine lakes (5,611 cells/mL). Hypereutrophic lakes had the highest phytoplankton cell concentration, with a mean concentration of 126,248 cells/mL. The mean cell concentration for eutrophic lakes was 86,354 cells/mL. Mesotrophic and oligotrophic lakes had the lowest mean cell concentrations of 51,269 and 18,793 cells/mL, respectively.

Blue-green algae (cyanobacteria) were most dominant in hypereutrophic lakes, accounting for 84.7 percent of all algal cells. Oligotrophic lakes had the lowest percent of blue-green algae present at 64.86 percent. Blue-green dominance had no notable variation for lake type. Natural lakes had the highest blue-green dominance of 87 percent, followed by 86 percent for impoundments and 84.5 percent for surface mine lakes.

*Secchi Disk Transparency and Trophic State*

Median Secchi depth for all lakes sampled was 1.7 meters (Figure 17). Impoundments has the lowest median Secchi depth of 1.05 meters, whereas surface mine lakes had a median Secchi depth of 3.1 meters. While natural lakes had a median Secchi depth of 1.70 meters, natural lakes represented the minimum and maximum Secchi depths of all lakes. Cedar Lake — in Lake County — has a Secchi depth of 0.15 meters, and Golden Lake — in Steuben County had a Secchi depth of 19 meters.




Figure 17 – Secchi depth distribution for 329 lakes sampled from 2015 to 2018.

Mesotrophic lakes were the most common based on TSI[chl-a], accounting for 44 percent of all lakes sampled (Figure 18). Eutrophic lakes accounted for 31 percent of lakes sampled. Only 9 percent of lakes sampled were oligotrophic. Airline Lake, in Greene County, had the lowest TSI[chl-a] of 22, whereas Waveland Lake, in Montgomery County, had a TSI[chl-a] of 79. Median TSI[chl-a] for all lakes sampled was 50.




Figure 18 – Number of lakes sampled from 2015 to 2018 (n = 329) by Carlson TSI [chl-a].

While TSI[chl-a] is reported consistently throughout this report, the relationship between TSI[chl-a] and TSI[TP] is important for the sampled lakes (Figure 19). Over half of the lakes sampled from 2015 to 2018 fall below the predicted relationship (red line in Figure 19) between chlorophyll-a and total phosphorus according to Carlson (1977).




Figure 19 – Carlson TSI [TP] plotted against Carlson TSI [chl-a] for 329 lakes sampled from 2015 to 2018. The red line indicated the predicted relationship between TSI [TP] and TSI [chl-a].

### Spatial Patterns

The 329 lakes sampled from 2015 to 2018 were located in 5 of the 6 ecoregions in Indiana (Figure 20). The Huron-Erie Lake Plain (Ecoregion 57) was the only ecoregion without a lake sampled. A majority of lakes sampled were in northeastern Indiana, with 58 percent occurring in the Southern Michigan/Northern Indiana Till Plain (Ecoregion 56).




Figure 20 – Location of lakes from 2015 to 2018 (n = 329) by Carlson TSI [chl-a] overlain on Indiana ecoregions (represented by color and corresponding number).

The Eastern Corn Belt Plains (Ecoregion 55) and the Central Corn Belt Plains (Ecoregion 54) had the highest median chlorophyll-a concentrations of 61.5 ug/L and 58 ug/L, respectively (Figure 21). The Interior River Lowland (Ecoregion 72) had the lowest median chlorophyll-a concentration of 43.5 ug/L.

Total phosphorus concentrations followed similar spatial patterns to that of chlorophyll-a concentrations (Figure 22). Ecoregion 54, 55, and 56 had higher median TP concentrations compared to that of Ecoregion 72 and 73. Median TP concentration was highest in Ecoregion 54 with a concentration of 0.123 mg/L, and Ecoregion 71 had the lowest median concentration of 0.039 mg/L.

Median Secchi depths were highest in Ecoregion 72 (2.55 meters) and 56 (1.70 meters) (Figure 23). Ecoregion 55 — the Eastern Corn Belt Plains — had the lowest median Secchi depth of 0.90 meters. While the Southern Michigan/Northern Indiana Till Plain (Ecoregion 56) had a median Secchi depth of 1.70 meters, the ecoregion also contained the lake with the highest Secchi depth (Golden Lake, Steuben Co.) of 19 meters.




Figure 21 – Chlorophyll-a (chl-a) distribution by ecoregion for 329 lakes sampled from 2015 to 2018. Figure (A) illustrates the total distribution by a dot plot and (B) illustrates the same distribution with a box plot.




Figure 22 – Total phosphorus (TP) distribution by ecoregion for 320 lakes sampled from 2015 to 2018 by the (A) total TP distribution and the (B) TP distribution under 0.50 mg/L.

TSI[chl-a] median values followed similar trends across ecoregions with chlorophyll-a and total phosphorus (Figure 24). Median TSI[chl-a] values were highest in Ecoregion 54 and 55, both of which were above the bottom limit of the eutrophic classification of 51. Ecoregion 72 had the lowest median TSI[chl-a] value of 43.5.




Figure 23 – Secchi depth distribution by ecoregion for 329 lakes sampled from 2015 to 2018 by the (A) total Secchi depth distribution and (B) Secchi depth values under 8 meters.




Figure 24 – Carlson TSI [chl-a] distribution by ecoregion for 329 lakes from 2015 to 2018. Figure (A) illustrates the total distribution by a dot plot and (B) illustrates the same distribution with a dox plot. The dashed line indicates the TSI break between eutrophic and mesotrophic.

## Lake Type Characteristics

Natural lakes represented the most common lake type sampled during the project, accounting for 58 percent of all lakes sampled. Impoundments, or reservoirs, represented 24 percent of lakes sampled, and 17 percent of lakes were surface mine lakes.

Impoundments had the highest median surface area of all lake types, but the lowest median maximum depth (Figure 25, Figure 26). Impoundments also had the largest variation in surface area from 0.81 to 4,353.75 hectares. While the median surface area of natural lakes was less than half of the median for impoundments, natural lakes were the deepest of the three lake types. The median surface area and maximum depth for natural lakes was 30.35 hectares and 10.70 meters, respectively. Median surface area for surface mine lakes was the smallest of the three lake types, with a value of 2.63 hectares. Surface mine lakes were generally deeper than that of impoundments but shallower compared to natural lakes, with a median maximum depth of 7.62 meters. The largest lake sampled was Lake Monroe (Monroe Co.) and Tippecanoe Lake (Kosciusko Co.) was the deepest lake sampled.




Figure 25 – Surface area distribution by lake type for 329 lakes sampled from 2015 to 2018 by the **(A)** total surface are distribution and **(B)** distribution under 850 hectares (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).




Figure 26 – Maximum depth distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure (A) illustrates the total distribution by a dot plot and (B) illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Natural lakes had the highest median average alkalinity concentration of 187.5 mg CaCO<sub>3</sub>/L (Figure 27). Median alkalinity concentration for surface mine lakes was 153 mg CaCO<sub>3</sub>/L, and surface mine lakes had the greatest alkalinity variation of the three lake types (47 to 1037.5 mg CaCO<sub>3</sub>/L). The median alkalinity concentration for impoundments was 115.5 mg CaCO<sub>3</sub>/L.




Figure 27 – Alkalinity distribution by lake type for 329 lakes sampled from 2015 to 2018 by (A) total alkalinity distribution and (B) distribution under 350 mg CaCO<sub>3</sub>/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Median average conductivity in surface mine lakes was almost twice the median of natural lakes, with conductivities of 872.50 and 457.83 umhos/cm, respectively (Figure 28). Surface mine lakes also had the greatest variation in conductivity, with values ranging from 158.15 to 2987.00 umhos/cm. Impoundments had the lowest conductivity of the lake types, with a median conductivity of 300.54 umhos/cm.




Figure 28 – Conductivity distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure (A) illustrates the total distribution by a dot plot and (B) illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Average pH values were consistent across the lake types, with only a 0.10 deviation between median pH values (Figure 29). Natural lakes had a median average pH of 7.6, and impoundments and surface mine lakes had the same median pH of 7.5.




Figure 29 – pH distribution by lake type for 329 lakes sampled from 2015 to 2018 by (A) total pH distribution and (B) distribution from 6 to 9 pH units (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Natural lakes had the highest median concentrations of NO<sub>3</sub>-N, NH<sub>3</sub>-N, and TP (Figure 30, Figure 31, Figure 32). Median concentrations of NO<sub>3</sub>-N and NH<sub>3</sub>-N in natural lakes were also more than twice the median concentration of impoundments and surface mine lakes. Median TP concentrations were less variable, with a variation of 0.01 mg/L between natural lakes and impoundments.




Figure 30 – Nitrate-nitrogen ( $\text{NO}_3\text{-N}$ ) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total  $\text{NO}_3\text{-N}$  distribution and **(B)** distribution under 0.80 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).




Figure 31 – Ammonia-nitrogen ( $\text{NH}_3\text{-N}$ ) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total  $\text{NH}_3\text{-N}$  distribution and **(B)** distribution under 2.50 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).




Figure 32 – Total phosphorus (TP) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total TP distribution and **(B)** distribution under 0.40 mg/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Chlorophyll-a concentrations were highest in impoundments, with a median chlorophyll-a concentration of 17.900 ug/L (Figure 33). The maximum chlorophyll-a concentration across all lakes sample of 146.596 was also an impoundment. Natural lakes had the second highest median chlorophyll-a concentration of 7.423 followed by surface mine lakes of 3.183.




Figure 33 – Chlorophyll-a (chl-a) distribution by lake type for 329 lakes sampled from 2015 to 2018 by **(A)** total chl-a distribution and **(B)** distribution under 80 ug/L (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

Secchi depths by lake type followed an inverse relationship to that of chlorophyll-a (Figure 34). Median Secchi depth for surface mine lakes was 3.10 meters, 1.70 meters for natural lakes, and 1.05 meters for impoundments.

Median TSI[chl-a] values for impoundments were greater than the bottom limit for the eutrophic interpretation, with a median of 59 (Figure 35). The median value for natural lakes was only 1 unit from the bottom of the eutrophic limit as well, with a value of 50. Median TSI[chl-a] for surface mine lakes was 42. Overall, 60 percent of impoundments were either eutrophic or hypereutrophic, compared to 46 percent of natural lakes and 14 percent of surface mine lakes.




Figure 34 – Secchi depth distribution by lake type for 329 lakes sampled from 2015 to 2018 by (A) total Secchi depth distribution and (B) distribution under 6 meters (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).




Figure 35 – TSI [chl-a] distribution by lake type for 329 lakes sampled from 2015 to 2018. Figure (A) illustrates the total TSI [chl-a] distribution by a dot plot and (B) illustrates the same distribution with a box plot (Impd = impoundments; NL = natural lakes; SML = surface mine lakes).

## DISCUSSION

### State of Indiana Lakes

Many lakes throughout the state of Indiana receive high nutrient loads, and thus are productive aquatic systems. This was expected as agricultural activity is a dominant land use throughout the state, and the subsequent runoff of nutrients — specifically nitrogen and phosphorus — would contribute to increased productivity in lakes.

While some lakes sampled had excessive levels of nitrogen, it appears *qualitatively* that many of the lakes sampled experience high phosphorus loading. Unlike nitrogen, phosphorus is bound in rock and sediment. As a result, phosphorus loading can occur at the water-sediment interface in the presence of anoxic conditions. Increased levels of SRP in the hypolimnion was common throughout the lakes sampled and indicates two interactions: the loading of phosphorus and the presence of thermal stratification. A molar relationship between nitrogen, phosphorus, and carbon — commonly referred to as the Redfield Ratio — is required to *quantitatively* determine if phosphorus is indeed the most common limiting nutrient for Indiana lakes.

Nitrogen and phosphorus are the primary nutrients for plant growth on lake and in the water. As a result, increased nutrient loading can contribute to increased algal growth. Algal communities occur in nearly all lakes and thus the presence of these communities is not necessarily an indication of impairment. However, certain algal groups are a concern due to their ability to produce nuisance blooms that can result in harmful conditions to both human and environmental health. Cyanobacteria (blue-green algae) are of particular concern in Indiana, and were common in many lakes sampled. Specifically, 27 percent of lakes sampled had blue-green communities that accounted for over 90 percent of all algal cells present. Understanding the presence of cyanobacteria cell density in lakes is an important management tool to identify the potential of harmful algal blooms (HABs) that can occur not only in the growing season, but throughout the entire year.

Trophic state is perhaps the most useful measure of the current state of Indiana Lakes, as well as a tool to compare Indiana to other states and regions across the United States. We found that nearly half (47 percent) of Indiana lakes were either eutrophic or hypereutrophic based on TSI [chl-a], indicating high levels of productivity. We did find some deviation in the relationship between the predicted relationship between TSI [chl-a] and TSI [TP]. According to Carlson (1977), chlorophyll-a concentrations can be predicted based on the TP concentration in the lake. However, we found that over half of the actual values were less than the predicted values. Non-algal turbidity is likely driving this deviation. Indiana Lakes are generally more turbid as a result of sediment runoff compared to the lakes that Carlson used in his model. Increased non-algal turbidity would reduce light penetration, decreasing the depth of the euphotic zone, and thus decrease algal photosynthesis. Therefore, by leveraging the known relationship with Carlson TSI values, we can gain additional insight on the function of Indiana lakes.

## **Spatial Patterns**

Aggregating lakes by ecoregion is helpful to identify region differences in lake water quality. Ecoregion 54 (Eastern Corn Belt Plains) and 55 (Central Corn Belt Plains) had higher median values for chlorophyll-a, TP, and TSI [chl-a] compared to Ecoregions 56, 71, and 72. Row crop agriculture is the primary land use with Ecoregions 54 and 55. The relationship between agricultural fertilizers and lake eutrophication is well-established, and the high relevance of agriculture in these ecoregion is likely the cause of increased nutrient and trophic state (Novotny 2003).

Ecoregion 71 and 72 are located in southern Indiana, and have less agricultural activity, more forested land, and are primarily impoundments. Even though reservoirs have larger watersheds and have increased potential for larger nutrient loads, these ecoregions all had lower median values. This finding indicates the importance of land use on lake water quality. Unsurprisingly, ecoregion 71 and 72 had higher median Secchi depth measurements compared to ecoregion 54 and 55.

While we see qualitative differences in our lakes aggregated by ecoregion, further statistical analysis is needed to develop a quantitative comparison. A previous study conducted by Tetra Tech (2008) concluded that there were no significant differences between the geographic regions of Indiana in terms of water quality. However, their analysis instead concluded that there were significant differences between the three dominant lake types in Indiana.

## **Lake Type Patterns**

Limnological characteristics can vary greatly with lake type. Our data included three lake types: natural lakes, impoundments, and surface mine lakes. Impoundments generally had a larger surface area, were shallower, and more productive compared to natural lakes and surface mine lakes. This finding was expected as larger watersheds can contribute higher nutrient loads and shallower lakes have a large portion of the water column in the euphotic zone, contributing to increased productivity.

Natural lakes were the deepest lakes sampled, and had median hypolimnetic nutrient samples (e.g. TP, SRP) that were higher than that of impoundments and surface mine lakes. This is likely from the depth of the lake promoting increased thermal stability, causing the hypolimnion to be anoxic longer into the growing season, and promoting the release of sediment-bound phosphorus into the lake.

Surface mine lakes were unique compared to the other lake types. Surface mine lakes had high median alkalinity concentrations and extremely high conductivity values compared to the other lake types. Most surface mine lakes are located in southwestern Indiana which is characterized by limestone geology. As a result, these lakes have higher concentrations of calcium carbonate (CaCO<sub>3</sub>). High conductivity in surface mine lakes is likely a byproduct of the mining process, where iron-sulfur compounds in mine waste can leach ions out of the soil and into the water

column (Gyure et al. 1987). As conductivity is a measure of the ability of water to pass an electrical current, increased concentrations of dissolved ions cause higher conductivities in these lakes.

## **CONCLUSIONS**

Summary conclusions from the 2015 to 2018 lake water quality assessment program include:

- Phosphorus concentrations in many Indiana lakes can be excessive and contribute to eutrophication.
- Internal phosphorus from lake sediments is an important source of phosphorus in many lakes, and is inherently difficult to control.
- High non-algal turbidity in many Indiana lakes results in reduced algal communities otherwise predicted by available phosphorus concentrations.
- Cyanobacteria (blue-green algae) are common in Indiana lakes, and should be monitored on lakes where cell dominance is high.
- Almost half of Indiana lakes assessed were either eutrophic or hypereutrophic. However, 44 percent of lakes were mesotrophic.
- Impoundments were generally the most productive lakes assessed.
- Carlson's Trophic State Index is a useful measure of overall trophic state in Indiana Lakes.
- Our randomized lake selection process — on average — generates data representative of Indiana lakes.

## REFERENCES

- Addy, Kelly, Linda Green and Elizabeth Herron. 2004. pH and Alkalinity. Watershed Watch, Cooperative Extension, University of Rhode Island, Kingston, Rhode Island.
- APHA. 2005. Standard Methods for the Examination of Water and Wastewater, 21st edition. American Public Health Association, Washington, D.C.
- Carlson, R.E. 1977. A trophic state index for lakes. *Limnology and Oceanography*, 2(2):361-369.
- CLP. 2011. Open Files, Indiana Clean Lakes Program, School of Public & Environmental Affairs, Indiana University, Bloomington, IN.
- Cooke, G.D. et al. 1993. *Management of Lakes and Reservoirs*, Second Edition. Lewis Publishers, Ann Arbor, Michigan.
- Gyure, R.A., A. Konopka, A. Brooks, and W. Doemel. 1987. Algal and Bacterial Activities in Acidic (pH 3) Strip Mine Lakes. *Applied and Environmental Microbiology*. 53(9): 2069-2076.
- Hutchinson, G.E. 1957. A Treatise on Limnology. *Volume I: Geography, Physics, and Chemistry*. John Wiley and Sons, Inc., New York.
- Jones, W. 1996. *Indiana lake Water Quality Update for 1989-1993*. Indiana Department of Environmental Management Clean Lakes Program, Indianapolis, Indiana.
- Novotny, V. 2003. *Water Quality: Diffuse Pollution and Watershed Management*, 2<sup>nd</sup> Edition. Wiley.
- Olem, H. and G. Flock, eds. 1990. *Lake and Reservoir Restoration Guidance Manual*, 2<sup>nd</sup> Edition. EPA 440/4-90-006. Prep. By N. Am. Lake Manage. Soc. For U.S. EPA, Washington, D.C.
- Omernik, J.M. and A.L. Gallant. 1988. *Ecoregions of the Upper Midwest*. EPA/600/3-88/037. U.S. Environmental Protection Agency, Environmental Research laboratory, Corvallis, Oregon.
- Prescott, G.W. 1982. *Algae of the Western Great Lakes Area*. Otto Koeltz Science Publishers, West Germany.
- PhycoTech. 2018. Business and Technical Information. PhycoTech, Inc., St. Joseph, MI
- St. Amand, Ann. 2010. Aquatic Microorganism Image Library (CD). PhycoTech, Inc., St. Joseph, MI

Tetra Tech. 2008. Data Analysis Report for Analysis of Indiana Lake Nutrient and Biological Data for the Nutrient Scientific Technical Exchange Partnership Support (N-STEPS). Tetra Tech, Inc., Owings Mills, MD.

US EPA. 1993. Methods for the Determination of Inorganic Substances in Environmental Samples; EPA-600/R-93/100. U.S. Environmental Protection Agency, Office of Research and Development, Cincinnati, OH.

Ward, H.B. and G.C. Whipple. 1959. Freshwater Biology, Second Edition. W.T. Edmondson, editor. John Wiley & Sons, Inc., New York.

Wehr, J.D. and R.G. Sheath. 2003. Freshwater Algae of North America, Ecology and Classification. Academic Press, San Diego.

Whitford, L.A. and G.J. Schumacher. 1984. A Manual of Fresh-Water Algae. Sparks Press, Raleigh, N.C.

## **APPENDICES**

Appendix A – Information for Indiana lakes sampled from 2015 to 2018.

| Lake Name              | County      | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------|-------------|-------------------|------|-------------------|---------------|
| Atwood                 | LaGrange    | impoundment       | 2015 | 7.28              | 10.1          |
| Bass                   | Sullivan    | Surface Mine Lake | 2015 | 85.39             | 16.0          |
| Bass (N.Chain)         | St. Joseph  | natural lake      | 2015 | 35.61             | 9.6           |
| Bear                   | Noble       | natural lake      | 2015 | 55.04             | 17.2          |
| Big Otter              | Steuben     | natural lake      | 2015 | 27.92             | 12.2          |
| Big Turkey             | LaGrange    | natural lake      | 2015 | 182.12            | 20.1          |
| Blackman               | LaGrange    | natural lake      | 2015 | 25.90             | 16.9          |
| Bobcat                 | Greene      | Surface Mine Lake | 2015 | 2.00              | 9.0           |
| Boones Pond            | Boone       | Surface Mine Lake | 2015 | 3.24              | 8.4           |
| Brush Creek Reservoir  | Jennings    | impoundment       | 2015 | 67.58             | 7.6           |
| Cagles Mill (Cataract) | Putnam      | impoundment       | 2015 | 566.58            | 12.5          |
| Canada                 | Porter      | natural lake      | 2015 | 4.05              | 7.0           |
| Carr                   | Kosciusko   | natural lake      | 2015 | 25.90             | 11.5          |
| Cedar                  | Lake        | natural lake      | 2015 | 316.07            | 4.2           |
| Center                 | Kosciusko   | natural lake      | 2015 | 48.56             | 13.0          |
| Clear (LaPorte)        | LaPorte     | natural lake      | 2015 | 42.90             | 3.6           |
| Crane                  | Noble       | natural lake      | 2015 | 11.33             | 10.9          |
| Crystal                | Greene      | Surface Mine Lake | 2015 | 3.24              | 11.0          |
| Failing                | Steuben     | natural lake      | 2015 | 9.31              | 12.0          |
| Fish (Lower)           | LaPorte     | natural lake      | 2015 | 54.23             | 4.7           |
| Fish (Upper)           | LaPorte     | natural lake      | 2015 | 56.25             | 7.1           |
| Gambill                | Sullivan    | Surface Mine Lake | 2015 | 4.86              | 13.0          |
| Geist Reservoir        | Marion      | impoundment       | 2015 | 728.46            | 7.2           |
| Goldeneye              | Kosciusko   | impoundment       | 2015 | 8.09              | 4.0           |
| Goodman                | Greene      | Surface Mine Lake | 2015 | 1.21              | 7.9           |
| Goose                  | Kosciusko   | natural lake      | 2015 | 10.93             | 12.2          |
| Green                  | Steuben     | natural lake      | 2015 | 9.71              | 10.7          |
| Griffy                 | Monroe      | impoundment       | 2015 | 52.61             | 9.0           |
| Grouse Ridge           | Bartholomew | impoundment       | 2015 | 8.09              | 8.6           |
| Hackberry              | Sullivan    | Surface Mine Lake | 2015 | 2.02              | 8.3           |
| Hale                   | Sullivan    | Surface Mine Lake | 2015 | 6.07              | 8.5           |
| Hartz                  | Starke      | natural lake      | 2015 | 11.33             | 10.1          |
| Hindman                | Noble       | natural lake      | 2015 | 5.26              | 6.0           |
| Hog                    | LaPorte     | natural lake      | 2015 | 23.88             | 16.8          |
| Huntingburg City       | Dubois      | impoundment       | 2015 | 73.25             | 6.1           |
| James                  | Kosciusko   | natural lake      | 2015 | 108.05            | 19.0          |
| John Hay               | Washington  | impoundment       | 2015 | 84.99             | 7.9           |
| Kickapoo               | Sullivan    | impoundment       | 2015 | 12.14             | 11.6          |
| King                   | Fulton      | natural lake      | 2015 | 7.69              | 9.8           |
| Larwill                | Whitley     | natural lake      | 2015 | 4.05              | 11.2          |

| Lake Name                    | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------------|------------|-------------------|------|-------------------|---------------|
| Little Bause                 | Noble      | natural lake      | 2015 | 2.83              | 5.0           |
| Long                         | Porter     | natural lake      | 2015 | 26.31             | 8.6           |
| Loon                         | Steuben    | natural lake      | 2015 | 55.85             | 18.0          |
| Loon                         | Whitley    | natural lake      | 2015 | 6.07              | 29.2          |
| Manitou                      | Fulton     | natural lake      | 2015 | 288.55            | 13.7          |
| Mansfield Reservoir (Hardin) | Parke      | impoundment       | 2015 | 833.68            | 18.5          |
| Marsh                        | Steuben    | natural lake      | 2015 | 22.66             | 12.0          |
| Martin                       | LaGrange   | natural lake      | 2015 | 10.52             | 17.5          |
| Mud (Chain of Lakes)         | Noble      | natural lake      | 2015 | 3.24              | 6.0           |
| Nauvoo                       | LaGrange   | natural lake      | 2015 | 15.38             | 9.0           |
| North Little                 | Kosciusko  | natural lake      | 2015 | 4.86              | 8.5           |
| Oliver                       | LaGrange   | natural lake      | 2015 | 150.14            | 27.0          |
| Oswego                       | Kosciusko  | natural lake      | 2015 | 16.59             | 11.0          |
| Otter                        | Steuben    | natural lake      | 2015 | 47.75             | 9.5           |
| Port Mitchell                | Noble      | natural lake      | 2015 | 6.07              | 9.9           |
| Prairie Creek Reservoir      | Delaware   | impoundment       | 2015 | 492.12            | 9.0           |
| Red Pine                     | Sullivan   | Surface Mine Lake | 2015 | 1.62              | 3.7           |
| Redbud                       | Sullivan   | Surface Mine Lake | 2015 | 1.62              | 7.0           |
| Ridinger                     | Kosciusko  | natural lake      | 2015 | 55.04             | 12.0          |
| Round                        | Whitley    | natural lake      | 2015 | 53.00             | 19.6          |
| Royer                        | LaGrange   | natural lake      | 2015 | 27.92             | 17.7          |
| Scales                       | Warrick    | Surface Mine Lake | 2015 | 26.71             | 6.3           |
| Schlamm                      | Clark      | impoundment       | 2015 | 285.31            | 6.9           |
| Shakamak                     | Sullivan   | impoundment       | 2015 | 22.66             | 6.6           |
| Shipshewana                  | LaGrange   | natural lake      | 2015 | 81.75             | 4.7           |
| Shriner                      | Whitley    | natural lake      | 2015 | 7.28              | 22.4          |
| Skunk                        | Greene     | Surface Mine Lake | 2015 | 0.30              | 5.0           |
| St. Joseph Reservoir         | Allen      | impoundment       | 2015 | 12.14             | 3.0           |
| Starve Hollow                | Jackson    | impoundment       | 2015 | 58.68             | 4.3           |
| Summit                       | Henry      | impoundment       | 2015 | 329.83            | 14.9          |
| Sycamore                     | Greene     | Surface Mine Lake | 2015 | 2.83              | 7.6           |
| T Lake                       | Sullivan   | Surface Mine Lake | 2015 | 2.02              | 7.6           |
| Todd                         | Greene     | Surface Mine Lake | 2015 | 3.24              | 10.0          |
| Trimble                      | Greene     | Surface Mine Lake | 2015 | 3.64              | 3.0           |
| Turtle                       | Sullivan   | Surface Mine Lake | 2015 | 3.24              | 7.0           |
| Waveland                     | Montgomery | impoundment       | 2015 | 145.69            | 5.0           |
| Wawasee                      | Kosciusko  | natural lake      | 2015 | 1059.50           | 22.8          |
| Webster                      | Kosciusko  | natural lake      | 2015 | 313.24            | 15.5          |
| White Oak #2                 | Knox       | impoundment       | 2015 | 2.83              | 5.2           |
| Willow                       | Sullivan   | Surface Mine Lake | 2015 | 1.62              | 9.8           |
| Yellowwood                   | Brown      | impoundment       | 2015 | 53.83             | 8.5           |

| Lake Name              | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------|------------|-------------------|------|-------------------|---------------|
| Atwood                 | LaGrange   | natural lake      | 2016 | 68.80             | 9.8           |
| Ball                   | Steuben    | natural lake      | 2016 | 35.21             | 19.8          |
| Banning                | Kosciusko  | natural lake      | 2016 | 4.86              | 4.6           |
| Bartley                | Noble      | natural lake      | 2016 | 13.76             | 9.3           |
| Bass                   | Starke     | natural lake      | 2016 | 544.32            | 6.7           |
| Bear                   | Noble      | natural lake      | 2016 | 55.04             | 16.7          |
| Beaver Dam             | Steuben    | natural lake      | 2016 | 4.45              | 7.2           |
| Bixler                 | Noble      | natural lake      | 2016 | 47.35             | 11.6          |
| Brush Creek Reservoir  | Jennings   | impoundment       | 2016 | 67.58             | 7.6           |
| Cagles Mill (Cataract) | Putnam     | impoundment       | 2016 | 566.58            | 11.7          |
| Cedar                  | Lake       | natural lake      | 2016 | 316.07            | 3.9           |
| Celina                 | Perry      | impoundment       | 2016 | 66.37             | 15.5          |
| Chapel pit             | Greene     | Surface Mine Lake | 2016 | 1.21              | 5.8           |
| Chrisney               | Spencer    | impoundment       | 2016 | 10.52             | 4.0           |
| Crane                  | Noble      | natural lake      | 2016 | 11.33             | 10.8          |
| Dallas                 | LaGrange   | natural lake      | 2016 | 114.53            | 30.2          |
| Diamond                | Noble      | natural lake      | 2016 | 42.49             | 23.2          |
| Eagle Creek Reservoir  | Marion     | impoundment       | 2016 | 611.10            | 13.1          |
| Elk Creek #9           | Washington | impoundment       | 2016 | 19.43             | 6.4           |
| Ferdinand City New     | Dubois     | impoundment       | 2016 | 4.05              | 4.3           |
| Ferdinand City Old     | Dubois     | impoundment       | 2016 | 6.07              | 5.5           |
| Fletcher               | Fulton     | natural lake      | 2016 | 18.21             | 12.0          |
| Freeman                | Carroll    | impoundment       | 2016 | 626.07            | 10.5          |
| Gage                   | Steuben    | natural lake      | 2016 | 132.34            | 22.0          |
| Gambill                | Sullivan   | Surface Mine Lake | 2016 | 4.86              | 12.8          |
| George                 | Steuben    | natural lake      | 2016 | 205.99            | 25.0          |
| George (Hobart)        | Lake       | impoundment       | 2016 | 109.27            | 2.1           |
| Gilbert                | Marshall   | natural lake      | 2016 | 14.16             | 8.8           |
| Goldeneye              | Kosciusko  | impoundment       | 2016 | 8.09              | 4.6           |
| Green Valley           | Vigo       | impoundment       | 2016 | 20.24             | 4.9           |
| Hackberry              | Sullivan   | Surface Mine Lake | 2016 | 2.02              | 9.0           |
| Hamilton               | Steuben    | natural lake      | 2016 | 324.57            | 21.3          |
| Hoffman                | Kosciusko  | natural lake      | 2016 | 75.68             | 9.4           |
| Holem                  | Marshall   | natural lake      | 2016 | 12.14             | 8.7           |
| Hunter                 | Elkhart    | natural lake      | 2016 | 40.07             | 8.5           |
| Knapp                  | Noble      | natural lake      | 2016 | 35.61             | 17.4          |
| Kuhn                   | Kosciusko  | natural lake      | 2016 | 47.75             | 8.1           |
| Lake of the Woods      | LaGrange   | natural lake      | 2016 | 55.04             | 25.7          |
| Lake of the Woods      | Marshall   | natural lake      | 2016 | 168.36            | 13.4          |
| Lemon                  | Monroe     | impoundment       | 2016 | 667.76            | 7.6           |
| Long                   | Porter     | natural lake      | 2016 | 26.31             | 9.1           |
| Lower Fry              | Sullivan   | Surface Mine Lake | 2016 | 1.62              | 2.1           |

| Lake Name               | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|-------------------------|------------|-------------------|------|-------------------|---------------|
| Maxinkuckee             | Marshall   | natural lake      | 2016 | 750.31            | 27.0          |
| Mayfield                | Sullivan   | Surface Mine Lake | 2016 | 6.07              | 7.6           |
| McClures                | Kosciusko  | natural lake      | 2016 | 12.95             | 8.7           |
| Miller (Chain-O)        | Noble      | natural lake      | 2016 | 4.45              | 8.8           |
| Molenkramer Reservoir   | Ripley     | impoundment       | 2016 | 37.64             | 2.1           |
| Monroe (Lower)          | Monroe     | impoundment       | 2016 | 4353.75           |               |
| Muncie                  | Noble      | natural lake      | 2016 | 19.02             | 8.2           |
| Nasby Mill Pond         | LaGrange   | impoundment       | 2016 | 14.16             | 1.4           |
| Nauvoo                  | LaGrange   | natural lake      | 2016 | 15.38             | 7.9           |
| North Twin              | LaGrange   | natural lake      | 2016 | 54.63             | 12.0          |
| Palestine               | Kosciusko  | impoundment       | 2016 | 93.89             | 8.2           |
| Prairie Creek Reservoir | Delaware   | impoundment       | 2016 | 492.12            | 8.8           |
| Pretty                  | LaGrange   | natural lake      | 2016 | 74.46             | 25.0          |
| Redbud                  | Sullivan   | Surface Mine Lake | 2016 | 1.62              | 7.0           |
| Rider                   | Noble      | natural lake      | 2016 | 2.02              | 4.9           |
| Robinson                | Whitley    | natural lake      | 2016 | 23.88             | 14.0          |
| Rothenberger            | Kosciusko  | natural lake      | 2016 | 2.43              | 14.6          |
| Sawmill                 | Kosciusko  | natural lake      | 2016 | 10.93             | 7.5           |
| Scheister               | Clay       | Surface Mine Lake | 2016 | 4.13              | 14.6          |
| Sellers                 | Kosciusko  | natural lake      | 2016 | 12.95             | 6.1           |
| Shaffer                 | White      | impoundment       | 2016 | 522.47            | 5.7           |
| Silver                  | Steuben    | natural lake      | 2016 | 96.32             | 10.5          |
| Simonton                | Elkhart    | natural lake      | 2016 | 114.13            | 7.3           |
| Skunk                   | Greene     | Surface Mine Lake | 2016 | 0.30              | 4.3           |
| South Twin              | LaGrange   | natural lake      | 2016 | 46.95             | 15.5          |
| Springs Valley (Tucker) | Orange     | impoundment       | 2016 | 57.06             | 9.1           |
| Spurgeon Hollow         | Washington | impoundment       | 2016 | 4.86              | 5.2           |
| Star                    | Greene     | Surface Mine Lake | 2016 | 2.02              | 7.6           |
| Steinbarger             | Noble      | natural lake      | 2016 | 29.54             | 11.3          |
| Still                   | LaGrange   | natural lake      | 2016 | 12.14             | 19.3          |
| Story (Lower)           | Dekalb     | natural lake      | 2016 | 31.16             | 9.1           |
| Story (Upper)           | Dekalb     | natural lake      | 2016 |                   | 8.7           |
| Sullivan                | Sullivan   | impoundment       | 2016 | 205.18            | 6.1           |
| Tippecanoe              | Kosciusko  | natural lake      | 2016 | 286.12            | 32.3          |
| University              | Monroe     | impoundment       | 2016 | 3.24              | 10.3          |
| Upper Long              | Noble      | natural lake      | 2016 | 34.80             | 14.0          |
| Webster                 | Kosciusko  | natural lake      | 2016 | 313.24            | 14.3          |
| Westler                 | LaGrange   | natural lake      | 2016 | 35.61             | 9.8           |
| Wolf                    | Lake       | natural lake      | 2016 | 155.81            | 4.7           |
| Adams                   | LaGrange   | natural lake      | 2017 | 118.58            | 27.7          |
| Airline                 | Greene     | Surface Mine Lake | 2017 | 10.12             | 21.0          |
| Atwood                  | LaGrange   | natural lake      | 2017 | 68.80             | 10.0          |

| Lake Name      | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|----------------|------------|-------------------|------|-------------------|---------------|
| Banning        | Kosciusko  | natural lake      | 2017 | 4.86              | 4.7           |
| Barton         | Steuben    | natural lake      | 2017 | 38.04             | 9.1           |
| Bass           | Starke     | natural lake      | 2017 | 544.32            | 7.0           |
| Benefiel       | Sullivan   | Surface Mine Lake | 2017 | 24.28             | 9.1           |
| Big Chapman    | Kosciusko  | natural lake      | 2017 | 167.55            | 13.0          |
| Big Turkey     | LaGrange   | natural lake      | 2017 | 182.12            | 19.4          |
| Bischoff Res.  | Ripley     | impoundment       | 2017 | 80.94             | 6.3           |
| Bowen          | Noble      | natural lake      | 2017 | 12.14             | 18.2          |
| Brookville     | Franklin   | impoundment       | 2017 | 2128.72           | 31.4          |
| Cedar          | LaGrange   | natural lake      | 2017 | 48.56             | 8.8           |
| Chapel Pit     | Greene     | Surface Mine Lake | 2017 | 1.21              | 6.0           |
| Chrisney       | Spencer    | impoundment       | 2017 | 10.52             | 4.3           |
| Clair          | Huntington | Surface Mine Lake | 2017 | 17.40             | 16.0          |
| Crooked        | Steuben    | natural lake      | 2017 | 324.57            | 18.6          |
| Crooked        | Whitley    | natural lake      | 2017 | 83.37             | 31.7          |
| Crosley        | Jennings   | impoundment       | 2017 | 5.67              | 6.1           |
| Dale Reservoir | Spencer    | impoundment       | 2017 | 13.36             | 5.8           |
| Deam           | Clark      | impoundment       | 2017 | 78.92             | 9.5           |
| Dock           | Noble      | natural lake      | 2017 | 6.48              | 6.7           |
| Eagle          | Noble      | natural lake      | 2017 | 32.78             | 13.1          |
| Engle          | Noble      | natural lake      | 2017 | 19.43             | 8.2           |
| Fish           | LaGrange   | natural lake      | 2017 | 40.47             | 24.8          |
| Fish           | Elkhart    | natural lake      | 2017 | 13.76             | 8.3           |
| Golden         | Steuben    | natural lake      | 2017 | 48.16             | 8.5           |
| Goldeneye      | Kosciusko  | impoundment       | 2017 | 8.09              | 4.8           |
| Griffy         | Monroe     | impoundment       | 2017 | 52.61             | 9.6           |
| Hale           | Sullivan   | Surface Mine Lake | 2017 | 6.07              | 9.1           |
| Hammond        | Greene     | Surface Mine Lake | 2017 | 2.43              | 8.0           |
| Hartz          | Starke     | natural lake      | 2017 | 11.33             | 10.0          |
| Henry          | Steuben    | natural lake      | 2017 | 8.09              | 6.2           |
| Horshoe        | Greene     | Surface Mine Lake | 2017 | 10.93             | 7.3           |
| Hunter         | Elkhart    | natural lake      | 2017 | 40.07             | 8.8           |
| Jimmerson      | Steuben    | natural lake      | 2017 | 114.53            | 16.8          |
| John Hay       | Washington | impoundment       | 2017 | 84.99             | 5.9           |
| Kiser          | Kosciusko  | natural lake      | 2017 | 3.64              | 6.0           |
| Latta          | Noble      | natural lake      | 2017 | 17.00             | 10.4          |
| Lawrence       | Marshall   | natural lake      | 2017 | 27.92             | 20.0          |
| Little Barbee  | Kosciusko  | natural lake      | 2017 | 27.52             | 7.1           |
| Long           | Porter     | natural lake      | 2017 | 26.31             | 8.2           |
| Loomis         | Porter     | natural lake      | 2017 | 25.09             | 16.7          |
| Loon           | Whitley    | natural lake      | 2017 | 6.07              | 28.4          |
| Manlove        | Fayette    | impoundment       | 2017 | 6.07              | 2.9           |

| Lake Name                    | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------------|------------|-------------------|------|-------------------|---------------|
| Mansfield Reservoir (Hardin) | Parke      | impoundment       | 2017 | 833.68            | 18.0          |
| Mayfield                     | Sullivan   | Surface Mine Lake | 2017 | 6.07              | 7.9           |
| McClish                      | Steuben    | natural lake      | 2017 | 14.16             | 16.9          |
| Mill Pond                    | Marshall   | natural lake      | 2017 | 52.20             | 5.5           |
| Miller (Chain-O)             | Noble      | natural lake      | 2017 | 4.45              | 7.3           |
| Mississinewa Reservoir       | Miami      | impoundment       | 2017 | 1286.95           | 21.6          |
| Molenkramer Res              | Ripley     | impoundment       | 2017 | 37.64             | 1.8           |
| Mongo Mill Pond              | LaGrange   | natural lake      | 2017 | 29.54             | 2.5           |
| Nasby Mill Pond              | LaGrange   | impoundment       | 2017 | 14.16             | 1.5           |
| Oak                          | Clark      | impoundment       | 2017 | 0.81              | 3.5           |
| Otter                        | Steuben    | natural lake      | 2017 | 47.75             | 8.8           |
| Patoka Reservoir             | Dubois     | impoundment       | 2017 | 3593.74           | 7.3           |
| Pleasant                     | Steuben    | natural lake      | 2017 | 171.59            | 15.0          |
| Price                        | Kosciusko  | natural lake      | 2017 | 4.86              | 13.0          |
| Riddles                      | St. Joseph | natural lake      | 2017 | 31.16             | 5.8           |
| Rider                        | Noble      | natural lake      | 2017 | 2.02              | 5.4           |
| Robinson                     | Whitley    | natural lake      | 2017 | 23.88             | 15.4          |
| Sawmill                      | Kosciusko  | natural lake      | 2017 | 10.93             | 7.7           |
| Schlamm                      | Clark      | impoundment       | 2017 | 7.69              | 6.6           |
| Simonton                     | Elkhart    | natural lake      | 2017 | 114.13            | 7.3           |
| Snow                         | Steuben    | natural lake      | 2017 | 125.46            | 24.7          |
| Starve Hollow                | Jackson    | impoundment       | 2017 | 58.68             | 4.8           |
| Steinbarger                  | Noble      | natural lake      | 2017 | 29.54             | 11.9          |
| Story (Upper)                | Dekalb     | natural lake      | 2017 |                   | 8.3           |
| Sylvan                       | Noble      | impoundment       | 2017 | 254.96            | 9.4           |
| Tipsaw                       | Perry      | impoundment       | 2017 | 574.00            | 6.0           |
| Trout                        | Sullivan   | Surface Mine Lake | 2017 | 2.02              | 6.0           |
| Upper Long                   | Noble      | natural lake      | 2017 | 34.80             | 16.2          |
| Versailles                   | Ripley     | impoundment       | 2017 | 93.08             | 4.2           |
| Waldron                      | Noble      | natural lake      | 2017 | 87.42             | 13.7          |
| Webster                      | Kosciusko  | natural lake      | 2017 | 313.24            | 14.5          |
| West                         | Sullivan   | Surface Mine Lake | 2017 | 39.20             | 25.0          |
| Westler                      | LaGrange   | natural lake      | 2017 | 35.61             | 10.0          |
| White Pine                   | Sullivan   | Surface Mine Lake | 2017 | 0.81              | 3.5           |
| Woods (Big Blue #3)          | Rush       | impoundment       | 2017 | 17.81             | 4.2           |
| Bass                         | Sullivan   | Surface Mine Lake | 2018 | 85.39             | 15.5          |
| Bass (N. Chain)              | St. Joseph | natural lake      | 2018 | 35.61             | 9.5           |
| Baughner                     | Noble      | natural lake      | 2018 | 12.95             | 10.7          |
| Big Chapman                  | Kosciusko  | natural lake      | 2018 | 167.55            | 12.0          |
| Big Fry                      | Sullivan   | Surface Mine Lake | 2018 | 1.82              | 3.3           |
| Big Long                     | LaGrange   | natural lake      | 2018 | 148.12            | 25.7          |
| Big Otter                    | Steuben    | natural lake      | 2018 | 27.92             | 12.7          |

| Lake Name              | County    | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------|-----------|-------------------|------|-------------------|---------------|
| Bobcat                 | Greene    | Surface Mine Lake | 2018 | 2.00              | 9.1           |
| Bowen                  | Noble     | natural lake      | 2018 | 12.14             | 18.8          |
| Brokesha               | LaGrange  | natural lake      | 2018 | 14.57             | 5.7           |
| Cagles Mill (Cataract) | Putnam    | impoundment       | 2018 | 566.58            | 10.7          |
| Canada                 | Porter    | natural lake      | 2018 | 4.05              | 6.7           |
| Chapel Pit             | Greene    | Surface Mine Lake | 2018 | 1.21              | 6.1           |
| Clear                  | Greene    | Surface Mine Lake | 2018 | 1.21              | 5.8           |
| Dallas                 | LaGrange  | natural lake      | 2018 | 114.53            | 31.0          |
| Duely                  | Noble     | natural lake      | 2018 | 8.50              | 5.8           |
| Engle                  | Noble     | natural lake      | 2018 | 19.43             | 8.1           |
| Failing "Gentian"      | Steuben   | natural lake      | 2018 | 9.31              | 10.0          |
| Ferdinand City New     | Dubois    | impoundment       | 2018 | 4.05              | 4.3           |
| Fish                   | Steuben   | natural lake      | 2018 | 23.88             | 8.1           |
| Fox                    | Sullivan  | Surface Mine Lake | 2018 |                   | 9.1           |
| Fry                    | Sullivan  | Surface Mine Lake | 2018 | 1.62              | 2.7           |
| George                 | Steuben   | natural lake      | 2018 | 205.99            | 24.9          |
| Golden                 | Steuben   | natural lake      | 2018 | 48.16             | 9.2           |
| Gooseneck              | Steuben   | natural lake      | 2018 | 10.12             | 10.3          |
| Gordy                  | Noble     | natural lake      | 2018 | 12.55             | 10.6          |
| Goshen Dam Pond        | Elkhart   | impoundment       | 2018 | 57.47             | 2.6           |
| Hogback                | Steuben   | natural lake      | 2018 | 59.09             | 7.9           |
| Impoundment No. 26     | Sullivan  | Surface Mine Lake | 2018 | 19.02             | 2.8           |
| J.C. Murphy            | Newton    | impoundment       | 2018 | 485.64            | 3.4           |
| Kings                  | Fulton    | natural lake      | 2018 | 7.69              | 9.5           |
| Knapp                  | Noble     | natural lake      | 2018 | 35.61             | 17.7          |
| Latta                  | Noble     | natural lake      | 2018 | 17.00             | 10.9          |
| Lawrence               | Marshall  | natural lake      | 2018 | 27.92             | 19.0          |
| Little Bause           | Noble     | natural lake      | 2018 | 2.83              | 4.8           |
| Little Chapman         | Kosciusko | natural lake      | 2018 | 48.56             | 9.2           |
| Little Otter           | Steuben   | natural lake      | 2018 | 13.76             | 10.5          |
| Little Pike            | Kosciusko | natural lake      | 2018 | 10.12             | 3.1           |
| Locust                 | Sullivan  | Surface Mine Lake | 2018 | 2.83              | 4.8           |
| Loomis                 | Porter    | natural lake      | 2018 | 25.09             | 16.5          |
| Loon                   | Steuben   | natural lake      | 2018 | 55.85             | 5.5           |
| McClish                | Steuben   | natural lake      | 2018 | 14.16             | 17.0          |
| Middlefork Res.        | Wayne     | impoundment       | 2018 | 112.10            | 9.7           |
| Mink                   | Porter    | natural lake      | 2018 | 14.16             |               |
| Mississinewa Res.      | Miami     | impoundment       | 2018 | 1286.95           | 18.6          |
| Nauvoo                 | LaGrange  | natural lake      | 2018 | 15.38             | 8.3           |
| Pike                   | Kosciusko | natural lake      | 2018 | 82.15             | 9.9           |
| Pleasant               | Steuben   | natural lake      | 2018 | 171.59            | 12.8          |
| Port Mitchell          | Noble     | natural lake      | 2018 | 6.07              | 9.7           |

| Lake Name              | County     | Lake Type         | Year | Surface Area (ha) | Max depth (m) |
|------------------------|------------|-------------------|------|-------------------|---------------|
| Potato Cr. (Worster)   | St. Joseph | impoundment       | 2018 | 132.34            | 5.4           |
| Prarie Creek Reservoir | Deleware   | impoundment       | 2018 | 492.12            | 8.8           |
| Pretty                 | LaGrange   | natural lake      | 2018 | 74.46             | 25.1          |
| Prides Creek           | Pie        | impoundment       | 2018 | 36.42             | 8.3           |
| Redbud                 | Sullivan   | Surface Mine Lake | 2018 | 1.62              |               |
| Riddles                | St. Joseph | natural lake      | 2018 | 31.16             | 5.6           |
| Robinson               | Whitley    | natural lake      | 2018 | 23.88             | 15.2          |
| Rothenberger           | Kosciusko  | natural lake      | 2018 | 2.43              | 8.5           |
| Sand                   | Noble      | natural lake      | 2018 | 19.02             | 15.9          |
| Scales                 | Warrick    | Surface Mine Lake | 2018 | 26.71             | 5.2           |
| Scheister              | Clay       | Surface Mine Lake | 2018 | 4.13              | 14.9          |
| Scott                  | Greene     | Surface Mine Lake | 2018 | 4.86              | 14.6          |
| Shakamak               | Sullivan   | impoundment       | 2018 | 22.66             | 6.7           |
| Silver                 | Steuben    | natural lake      | 2018 | 96.32             | 9.1           |
| Skunk                  | Greene     | Surface Mine Lake | 2018 | 0.30              | 4.5           |
| Spencer                | Sullivan   | Surface Mine Lake | 2018 | 2.43              | 5.5           |
| Starve Hollow          | Jackson    | impoundment       | 2018 | 58.68             | 4.9           |
| Steinbarger            | Noble      | natural lake      | 2018 | 29.54             | 12.0          |
| Stump Jumper           | Clay       | Surface Mine Lake | 2018 | 2.39              | 9.1           |
| Syl-Van                | Steuben    | natural lake      | 2018 | 9.71              | 11.0          |
| Syracuse               | Kosciusko  | natural lake      | 2018 | 228.25            | 10.1          |
| Tamarack               | LaPorte    | natural lake      | 2018 | 8.09              | 1.5           |
| Thomas                 | Marshall   | natural lake      | 2018 | 6.48              | 13.7          |
| Tippecanoe             | Kosciusko  | natural lake      | 2018 | 286.12            | 37.0          |
| Twin Pitts, East       | Pike       | Surface Mine Lake | 2018 | 12.55             | 5.0           |
| Twin Pitts, West       | Pike       | Surface Mine Lake | 2018 | 7.28              | 2.7           |
| University             | Monroe     | impoundment       | 2018 | 3.24              | 10.4          |
| Upper Long             | Noble      | natural lake      | 2018 | 34.80             | 16.3          |
| Versailles             | Ripley     | impoundment       | 2018 | 93.08             | 4.9           |
| Waveland               | Montgomery | impoundment       | 2018 | 145.69            | 6.0           |
| Webster                | Kosciusko  | natural lake      | 2018 | 313.24            | 15.0          |
| Woods (Big Blue #3)    | Rush       | impoundment       | 2018 | 17.81             | 4.3           |

Appendix B – Trophic state indices for all lakes sampled from 2015 to 2018.

| Lake Name              | County      | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|------------------------|-------------|------|----------|-------------|----------|
| Atwood                 | LaGrange    | 2015 | 52       | 46          | 46       |
| Bass                   | Sullivan    | 2015 | 39       | 40          | 49       |
| Bass (N.Chain)         | St. Joseph  | 2015 | 49       | 43          | 54       |
| Bear                   | Noble       | 2015 | 57       | 62          | 61       |
| Big Otter              | Steuben     | 2015 | 53       | 52          | 48       |
| Big Turkey             | LaGrange    | 2015 | 51       | 50          | 46       |
| Blackman               | LaGrange    | 2015 | 49       | 42          | 51       |
| Bobcat                 | Greene      | 2015 | 46       | 40          | 47       |
| Boones Pond            | Boone       | 2015 | 44       | 33          | 48       |
| Brush Creek Reservoir  | Jennings    | 2015 | 67       |             | 65       |
| Cagles Mill (Cataract) | Putnam      | 2015 | 67       | 64          |          |
| Canada                 | Porter      | 2015 | 55       | 54          | 51       |
| Carr                   | Kosciusko   | 2015 | 67       | 69          | 44       |
| Cedar                  | Lake        | 2015 | 83       | 77          | 74       |
| Center                 | Kosciusko   | 2015 | 46       | 47          | 55       |
| Clear (LaPorte)        | LaPorte     | 2015 | 60       | 51          | 57       |
| Crane                  | Noble       | 2015 | 56       | 60          | 68       |
| Crystal                | Greene      | 2015 | 52       |             | 59       |
| Failing                | Steuben     | 2015 | 47       | 42          | 44       |
| Fish (Lower)           | LaPorte     | 2015 | 62       | 51          | 51       |
| Fish (Upper)           | LaPorte     | 2015 | 57       | 58          | 56       |
| Gambill                | Sullivan    | 2015 | 29       | 33          | 41       |
| Geist Reservoir        | Marion      | 2015 | 60       | 61          | 64       |
| Goldeneye              | Kosciusko   | 2015 | 49       | 50          | 77       |
| Goodman                | Greene      | 2015 | 44       | 38          | 37       |
| Goose                  | Kosciusko   | 2015 | 45       | 42          | 41       |
| Green                  | Steuben     | 2015 | 56       | 35          | 42       |
| Griffy                 | Monroe      | 2015 | 36       | 33          | 37       |
| Grouse Ridge           | Bartholomew | 2015 | 55       | 55          | 51       |
| Hackberry              | Sullivan    | 2015 | 42       | 51          | 57       |
| Hale                   | Sullivan    | 2015 | 43       | 44          | 60       |
| Hartz                  | Starke      | 2015 | 56       | 53          | 53       |
| Hindman                | Noble       | 2015 | 52       | 57          | 55       |
| Hog                    | LaPorte     | 2015 | 47       | 45          | 44       |
| Huntingburg City       | Dubois      | 2015 | 67       |             | 58       |

| Lake Name                       | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|---------------------------------|------------|------|----------|-------------|----------|
| James                           | Kosciusko  | 2015 |          | 53          | 57       |
| John Hay                        | Washington | 2015 | 45       |             | 57       |
| Kickapoo                        | Sullivan   | 2015 | 38       |             | 52       |
| King                            | Fulton     | 2015 | 73       | 70          | 72       |
| Larwill                         | Whitley    | 2015 | 62       | 75          | 78       |
| Little Bause                    | Noble      | 2015 | 50       | 77          | 65       |
| Long                            | Porter     | 2015 | 49       | 47          | 51       |
| Loon                            | Steuben    | 2015 | 34       | 51          | 52       |
| Loon                            | Whitley    | 2015 | 59       | 73          | 63       |
| Manitou                         | Fulton     | 2015 | 73       | 61          |          |
| Mansfield Reservoir<br>(Hardin) | Parke      | 2015 | 63       | 65          |          |
| Marsh                           | Steuben    | 2015 | 50       | 49          | 47       |
| Martin                          | LaGrange   | 2015 | 55       | 60          | 65       |
| Mud (Chain of Lakes)            | Noble      | 2015 | 59       | 70          | 59       |
| Nauvoo                          | LaGrange   | 2015 | 62       | 58          | 61       |
| North Little                    | Kosciusko  | 2015 | 55       | 54          | 66       |
| Oliver                          | LaGrange   | 2015 | 52       | 48          | 54       |
| Oswego                          | Kosciusko  | 2015 | 51       | 43          | 58       |
| Otter                           | Steuben    | 2015 | 50       | 49          | 54       |
| Port Mitchell                   | Noble      | 2015 | 59       | 69          | 65       |
| Prairie Creek Reservoir         | Delaware   | 2015 | 57       | 59          | 71       |
| Red Pine                        | Sullivan   | 2015 | 52       | 64          | 67       |
| Redbud                          | Sullivan   | 2015 | 53       | 49          | 54       |
| Ridinger                        | Kosciusko  | 2015 | 70       | 65          | 57       |
| Round                           | Whitley    | 2015 | 54       | 57          | 52       |
| Royer                           | LaGrange   | 2015 | 63       | 62          | 54       |
| Scales                          | Warrick    | 2015 | 63       |             | 59       |
| Schlamm                         | Clark      | 2015 | 59       |             | 54       |
| Shakamak                        | Sullivan   | 2015 |          |             | 65       |
| Shipshewana                     | LaGrange   | 2015 | 77       | 70          | 75       |
| Shriner                         | Whitley    | 2015 | 47       | 57          | 57       |
| Skunk                           | Greene     | 2015 | 49       | 45          | 54       |
| St. Joseph Reservoir            | Allen      | 2015 | 75       | 65          | 78       |
| Starve Hollow                   | Jackson    | 2015 | 63       |             | 69       |
| Summit                          | Henry      | 2015 | 34       | 35          |          |

| Lake Name              | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|------------------------|------------|------|----------|-------------|----------|
| Sycamore               | Greene     | 2015 | 60       | 45          | 37       |
| T Lake                 | Sullivan   | 2015 | 47       | 39          | 49       |
| Todd                   | Greene     | 2015 | 35       | 35          | 37       |
| Trimble                | Greene     | 2015 | 54       | 46          | 47       |
| Turtle                 | Sullivan   | 2015 | 42       | 35          | 47       |
| Waveland               | Montgomery | 2015 | 77       | 75          |          |
| Wawasee                | Kosciusko  | 2015 | 47       | 40          | 53       |
| Webster                | Kosciusko  | 2015 | 52       | 43          | 58       |
| White Oak #2           | Knox       | 2015 | 65       |             | 65       |
| Willow                 | Sullivan   | 2015 | 42       |             | 44       |
| Yellowwood             | Brown      | 2015 | 47       | 41          | 53       |
| Atwood                 | LaGrange   | 2016 | 53       | 40          | 47       |
| Ball                   | Steuben    | 2016 | 61       | 56          | 40       |
| Banning                | Kosciusko  | 2016 | 52       | 54          | 53       |
| Bartley                | Noble      | 2016 | 60       | 63          | 57       |
| Bass                   | Starke     | 2016 | 83       | 71          | 70       |
| Bear                   | Noble      | 2016 | 52       | 57          | 58       |
| Beaver Dam             | Steuben    | 2016 | 46       | 42          | 51       |
| Bixler                 | Noble      | 2016 | 45       | 48          | 49       |
| Brush Creek Reservoir  | Jennings   | 2016 | 63       | 64          | 56       |
| Cagles Mill (Cataract) | Putnam     | 2016 | 73       | 71          | 60       |
| Cedar                  | Lake       | 2016 | 87       | 71          | 79       |
| Celina                 | Perry      | 2016 | 43       | 34          | 42       |
| Chapel pit             | Greene     | 2016 | 45       | 40          | 49       |
| Chrisney               | Spencer    | 2016 | 56       | 39          | 50       |
| Crane                  | Noble      | 2016 | 66       | 65          | 62       |
| Dallas                 | LaGrange   | 2016 | 59       | 48          | 48       |
| Diamond                | Noble      | 2016 | 52       | 50          | 41       |
| Eagle Creek Reservoir  | Marion     | 2016 | 57       | 59          | 57       |
| Elk Creek #9           | Washington | 2016 | 44       | 46          | 47       |
| Ferdinand City New     | Dubois     | 2016 | 73       |             | 82       |
| Ferdinand City Old     | Dubois     | 2016 | 53       | 61          | 53       |
| Fletcher               | Fulton     | 2016 | 48       | 47          | 47       |
| Freeman                | Carroll    | 2016 | 60       | 64          | 63       |
| Gage                   | Steuben    | 2016 | 43       | 30          | 36       |

| Lake Name               | County    | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|-------------------------|-----------|------|----------|-------------|----------|
| Gambill                 | Sullivan  | 2016 | 29       | 27          | 44       |
| George                  | Steuben   | 2016 | 32       | 33          | 40       |
| George (Hobart)         | Lake      | 2016 | 69       | 70          | 75       |
| Gilbert                 | Marshall  | 2016 | 64       | 62          | 58       |
| Goldeneye               | Kosciusko | 2016 | 49       | 47          | 47       |
| Green Valley            | Vigo      | 2016 | 69       | 70          | 69       |
| Hackberry               | Sullivan  | 2016 | 37       | 39          | 40       |
| Hamilton                | Steuben   | 2016 | 45       | 44          | 46       |
| Hoffman                 | Kosciusko | 2016 | 55       | 52          | 53       |
| Holem                   | Marshall  | 2016 | 47       | 44          | 49       |
| Hunter                  | Elkhart   | 2016 | 44       | 43          | 52       |
| Knapp                   | Noble     | 2016 | 50       | 44          | 44       |
| Kuhn                    | Kosciusko | 2016 | 44       | 39          | 45       |
| Lake of the Woods       | LaGrange  | 2016 | 41       | 41          | 54       |
| Lake of the Woods       | Marshall  | 2016 | 49       | 55          | 42       |
| Lemon                   | Monroe    | 2016 | 70       | 68          | 57       |
| Long                    | Porter    | 2016 | 49       | 47          | 53       |
| Lower Fry               | Sullivan  | 2016 | 52       | 52          | 44       |
| Maxinkuckee             | Marshall  | 2016 | 58       | 41          | 40       |
| Mayfield                | Sullivan  | 2016 | 36       | 34          | 32       |
| McClures                | Kosciusko | 2016 | 67       | 69          | 65       |
| Miller (Chain-O)        | Noble     | 2016 | 52       | 56          | 56       |
| Molenkramer Reservoir   | Ripley    | 2016 | 77       | 76          | 83       |
| Monroe (Lower)          | Monroe    | 2016 |          | 49          | 20       |
| Muncie                  | Noble     | 2016 | 64       | 70          | 66       |
| Nasby Mill Pond         | LaGrange  | 2016 | 54       | 30          | 51       |
| Nauvoo                  | LaGrange  | 2016 | 52       | 50          | 47       |
| North Twin              | LaGrange  | 2016 | 47       | 40          | 32       |
| Palestine               | Kosciusko | 2016 | 66       | 69          | 74       |
| Prairie Creek Reservoir | Delaware  | 2016 | 57       | 56          | 48       |
| Pretty                  | LaGrange  | 2016 | 40       | 36          | 36       |
| Redbud                  | Sullivan  | 2016 | 41       | 38          | 40       |
| Rider                   | Noble     | 2016 | 54       | 54          | 49       |
| Robinson                | Whitley   | 2016 | 59       | 55          | 62       |
| Rothenberger            | Kosciusko | 2016 | 41       | 43          | 45       |

| Lake Name               | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|-------------------------|------------|------|----------|-------------|----------|
| Sawmill                 | Kosciusko  | 2016 | 63       | 60          | 58       |
| Scheister               | Clay       | 2016 | 38       | 39          | 40       |
| Sellers                 | Kosciusko  | 2016 | 67       | 70          | 71       |
| Shaffer                 | White      | 2016 | 62       | 64          | 66       |
| Silver                  | Steuben    | 2016 | 42       | 39          | 44       |
| Simonton                | Elkhart    | 2016 | 52       | 49          | 40       |
| Skunk                   | Greene     | 2016 | 49       | 50          | 49       |
| South Twin              | LaGrange   | 2016 | 52       | 39          | 32       |
| Springs Valley (Tucker) | Orange     | 2016 | 41       | 40          | 30       |
| Spurgeon Hollow         | Washington | 2016 | 55       | 55          | 54       |
| Star                    | Greene     | 2016 | 42       | 38          | 41       |
| Steinbarger             | Noble      | 2016 | 58       | 57          | 46       |
| Still                   | LaGrange   | 2016 | 56       | 48          | 45       |
| Story (Lower)           | Dekalb     | 2016 | 44       | 42          | 43       |
| Story (Upper)           | Dekalb     | 2016 | 53       | 52          | 44       |
| Sullivan                | Sullivan   | 2016 | 72       | 72          | 66       |
| Tippecanoe              | Kosciusko  | 2016 | 53       | 46          | 47       |
| University              | Monroe     | 2016 | 59       | 58          | 58       |
| Upper Long              | Noble      | 2016 | 52       | 49          | 54       |
| Webster                 | Kosciusko  | 2016 | 56       | 56          | 58       |
| Westler                 | LaGrange   | 2016 | 61       | 55          | 49       |
| Wolf                    | Lake       | 2016 | 61       | 65          | 65       |
| Adams                   | LaGrange   | 2017 | 49       | 47          | 39       |
| Airline                 | Greene     | 2017 | 33       | 22          | 14       |
| Atwood                  | LaGrange   | 2017 | 47       | 46          | 32       |
| Banning                 | Kosciusko  | 2017 | 52       |             | 57       |
| Barton                  | Steuben    | 2017 | 50       | 40          | 47       |
| Bass                    | Starke     | 2017 | 83       | 76          | 72       |
| Benefiel                | Sullivan   | 2017 | 34       |             | 14       |
| Big Chapman             | Kosciusko  | 2017 | 51       | 49          | 43       |
| Big Turkey              | LaGrange   | 2017 | 47       | 53          | 46       |
| Bischoff Res.           | Ripley     | 2017 | 67       | 63          | 68       |
| Bowen                   | Noble      | 2017 | 50       | 46          | 58       |
| Brookville              | Franklin   | 2017 | 46       | 52          | 44       |
| Cedar                   | LaGrange   | 2017 | 42       | 40          | 46       |

| Lake Name                       | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|---------------------------------|------------|------|----------|-------------|----------|
| Chapel Pit                      | Greene     | 2017 | 51       | 49          | 41       |
| Chrisney                        | Spencer    | 2017 | 46       | 32          | 30       |
| Clair                           | Huntington | 2017 | 62       | 37          | 69       |
| Crooked                         | Steuben    | 2017 | 49       | 37          | 47       |
| Crooked                         | Whitley    | 2017 | 39       | 39          | 55       |
| Crosley                         | Jennings   | 2017 | 59       | 58          | 63       |
| Dale Reservoir                  | Spencer    | 2017 | 70       | 69          | 70       |
| Deam                            | Clark      | 2017 | 51       | 28          | 47       |
| Dock                            | Noble      | 2017 | 60       | 61          | 66       |
| Eagle                           | Noble      | 2017 | 45       | 27          | 50       |
| Engle                           | Noble      | 2017 | 47       | 48          | 49       |
| Fish                            | LaGrange   | 2017 | 73       | 65          | 57       |
| Fish                            | Elkhart    | 2017 | 57       |             | 67       |
| Golden                          | Steuben    | 2017 | 18       | 54          | 54       |
| Goldeneye                       | Kosciusko  | 2017 | 46       | 41          | 54       |
| Griffy                          | Monroe     | 2017 | 40       | 35          | 46       |
| Hale                            | Sullivan   | 2017 | 40       | 28          | 20       |
| Hammond                         | Greene     | 2017 | 35       | 34          | 40       |
| Hartz                           | Starke     | 2017 | 46       | 50          | 46       |
| Henry                           | Steuben    | 2017 | 49       | 49          | 61       |
| Horshoe                         | Greene     | 2017 | 42       | 40          | 44       |
| Hunter                          | Elkhart    | 2017 | 48       |             | 32       |
| Jimmerson                       | Steuben    | 2017 | 48       | 43          | 56       |
| John Hay                        | Washington | 2017 | 40       | 37          | 51       |
| Kiser                           | Kosciusko  | 2017 | 47       | 47          | 53       |
| Latta                           | Noble      | 2017 | 55       | 42          | 40       |
| Lawrence                        | Marshall   | 2017 | 45       | 41          | 47       |
| Little Barbee                   | Kosciusko  | 2017 | 59       |             | 54       |
| Long                            | Porter     | 2017 | 47       | 43          | 48       |
| Loomis                          | Porter     | 2017 | 53       | 43          | 97       |
| Loon                            | Whitley    | 2017 | 62       | 65          | 62       |
| Manlove                         | Fayette    | 2017 | 73       | 66          | 80       |
| Mansfield Reservoir<br>(Hardin) | Parke      | 2017 | 63       | 50          | 54       |
| Mayfield                        | Sullivan   | 2017 | 38       | 35          | 42       |
| McClish                         | Steuben    | 2017 | 49       | 33          | 40       |

| Lake Name              | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|------------------------|------------|------|----------|-------------|----------|
| Mill Pond              | Marshall   | 2017 | 53       | 49          | 51       |
| Miller (Chain-O)       | Noble      | 2017 | 59       | 61          | 66       |
| Mississinewa Reservoir | Miami      | 2017 | 59       | 60          | 62       |
| Molenkramer Res        | Ripley     | 2017 | 77       | 62          | 80       |
| Mongo Mill Pond        | LaGrange   | 2017 | 57       | 44          | 68       |
| Nasby Mill Pond        | LaGrange   | 2017 | 56       | 63          | 60       |
| Oak                    | Clark      | 2017 | 65       | 58          | 59       |
| Otter                  | Steuben    | 2017 | 54       | 49          | 52       |
| Patoka Reservoir       | Dubois     | 2017 | 57       | 37          | 24       |
| Pleasant               | Steuben    | 2017 | 52       | 49          | 55       |
| Price                  | Kosciusko  | 2017 | 44       | 32          | 45       |
| Riddles                | St. Joseph | 2017 | 67       | 67          | 71       |
| Rider                  | Noble      | 2017 | 52       | 55          | 44       |
| Robinson               | Whitley    | 2017 | 67       | 71          | 64       |
| Sawmill                | Kosciusko  | 2017 | 59       | 62          | 57       |
| Schlamm                | Clark      | 2017 | 55       | 45          | 54       |
| Simonton               | Elkhart    | 2017 | 52       |             | 39       |
| Snow                   | Steuben    | 2017 | 47       | 46          | 51       |
| Starve Hollow          | Jackson    | 2017 | 56       | 51          | 56       |
| Steinbarger            | Noble      | 2017 | 59       | 61          | 67       |
| Story (Upper)          | Dekalb     | 2017 | 52       | 57          | 57       |
| Sylvan                 | Noble      | 2017 | 57       | 58          | 61       |
| Tipsaw                 | Perry      | 2017 | 56       | 48          | 47       |
| Trout                  | Sullivan   | 2017 | 52       | 52          | 57       |
| Upper Long             | Noble      | 2017 | 59       | 41          | 85       |
| Versailles             | Ripley     | 2017 | 77       | 62          | 74       |
| Waldron                | Noble      | 2017 | 59       | 65          | 64       |
| Webster                | Kosciusko  | 2017 | 49       | 54          | 55       |
| West                   | Sullivan   | 2017 | 60       | 43          | 52       |
| Westler                | LaGrange   | 2017 | 46       | 50          | 60       |
| White Pine             | Sullivan   | 2017 | 44       | 38          | 54       |
| Woods (Big Blue #3)    | Rush       | 2017 | 70       | 63          | 64       |
| Bass                   | Sullivan   | 2018 | 40       | 37          |          |
| Bass (N. Chain)        | St. Joseph | 2018 | 46       | 40          | 82       |
| Baughner               | Noble      | 2018 | 62       | 65          | 66       |

| Lake Name              | County    | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|------------------------|-----------|------|----------|-------------|----------|
| Big Chapman            | Kosciusko | 2018 | 50       | 47          | 53       |
| Big Fry                | Sullivan  | 2018 | 60       | 65          | 44       |
| Big Long               | LaGrange  | 2018 | 42       | 44          | 49       |
| Big Otter              | Steuben   | 2018 | 46       | 47          | 51       |
| Bobcat                 | Greene    | 2018 | 52       | 44          | 48       |
| Bowen                  | Noble     | 2018 | 67       | 72          | 40       |
| Brokesha               | LaGrange  | 2018 | 43       | 48          | 40       |
| Cagles Mill (Cataract) | Putnam    | 2018 | 62       | 65          | 52       |
| Canada                 | Porter    | 2018 | 33       | 50          |          |
| Chapel Pit             | Greene    | 2018 | 48       | 42          | 77       |
| Clear                  | Greene    | 2018 | 44       | 46          | 49       |
| Dallas                 | LaGrange  | 2018 | 54       | 45          | 46       |
| Duely                  | Noble     | 2018 | 49       | 56          | 43       |
| Engle                  | Noble     | 2018 | 50       | 51          | 59       |
| Failing "Gentian"      | Steuben   | 2018 | 44       | 35          | 45       |
| Ferdinand City New     | Dubois    | 2018 | 69       | 74          | 60       |
| Fish                   | Steuben   | 2018 | 57       | 68          | 51       |
| Fox                    | Sullivan  | 2018 | 59       | 51          | 52       |
| Fry                    | Sullivan  | 2018 | 49       | 48          | 39       |
| George                 | Steuben   | 2018 | 38       | 44          | 61       |
| Golden                 | Steuben   | 2018 | 60       | 58          | 66       |
| Gooseneck              | Steuben   | 2018 | 44       | 35          | 49       |
| Gordy                  | Noble     | 2018 | 43       | 47          |          |
| Goshen Dam Pond        | Elkhart   | 2018 | 63       | 42          | 51       |
| Hogback                | Steuben   | 2018 | 57       | 58          | 70       |
| Impoundment No. 26     | Sullivan  | 2018 | 67       | 69          | 56       |
| J.C. Murphy            | Newton    | 2018 | 67       | 70          | 43       |
| Kings                  | Fulton    | 2018 | 57       | 56          | 53       |
| Knapp                  | Noble     | 2018 | 49       | 51          | 63       |
| Latta                  | Noble     | 2018 | 57       | 45          | 74       |
| Lawrence               | Marshall  | 2018 | 43       | 37          | 72       |
| Little Bause           | Noble     | 2018 | 42       | 40          | 51       |
| Little Chapman         | Kosciusko | 2018 | 57       | 63          | 63       |
| Little Otter           | Steuben   | 2018 | 49       | 52          | 67       |
| Little Pike            | Kosciusko | 2018 | 63       | 67          | 45       |

| Lake Name              | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|------------------------|------------|------|----------|-------------|----------|
| Locust                 | Sullivan   | 2018 | 43       | 44          | 64       |
| Loomis                 | Porter     | 2018 | 51       | 52          | 80       |
| Loon                   | Steuben    | 2018 | 51       | 48          | 45       |
| McClish                | Steuben    | 2018 | 43       | 41          | 66       |
| Middlefork Res.        | Wayne      | 2018 | 70       | 63          | 57       |
| Mink                   | Porter     | 2018 |          |             | 55       |
| Mississinewa Res.      | Miami      | 2018 | 57       | 53          | 78       |
| Nauvoo                 | LaGrange   | 2018 | 56       | 58          | 44       |
| Pike                   | Kosciusko  | 2018 | 62       | 68          | 65       |
| Pleasant               | Steuben    | 2018 | 50       | 43          | 47       |
| Port Mitchell          | Noble      | 2018 | 55       | 57          | 50       |
| Potato Cr. (Worster)   | St. Joseph | 2018 | 65       | 63          | 51       |
| Prarie Creek Reservoir | Deleware   | 2018 | 57       | 61          | 57       |
| Pretty                 | LaGrange   | 2018 | 39       | 41          | 51       |
| Prides Creek           | Pie        | 2018 | 57       | 57          | 44       |
| Redbud                 | Sullivan   | 2018 |          | 44          | 47       |
| Riddles                | St. Joseph | 2018 | 64       | 69          | 62       |
| Robinson               | Whitley    | 2018 | 65       |             | 58       |
| Rothenberger           | Kosciusko  | 2018 | 49       | 53          | 54       |
| Sand                   | Noble      | 2018 | 54       | 62          | 65       |
| Scales                 | Warrick    | 2018 | 55       | 59          | 69       |
| Scheister              | Clay       | 2018 | 37       | 42          | 54       |
| Scott                  | Greene     | 2018 | 39       | 35          | 58       |
| Shakamak               | Sullivan   | 2018 | 60       | 62          | 45       |
| Silver                 | Steuben    | 2018 | 49       | 46          | 68       |
| Skunk                  | Greene     | 2018 | 48       | 46          | 49       |
| Spencer                | Sullivan   | 2018 | 47       | 46          | 53       |
| Starve Hollow          | Jackson    | 2018 | 60       | 63          | 74       |
| Steinbarger            | Noble      | 2018 | 66       | 71          | 46       |
| Stump Jumper           | Clay       | 2018 | 43       | 50          | 48       |
| Syl-Van                | Steuben    | 2018 | 46       | 43          | 55       |
| Syracuse               | Kosciusko  | 2018 | 47       | 50          | 56       |
| Tamarack               | LaPorte    | 2018 | 46       | 76          | 51       |
| Thomas                 | Marshall   | 2018 | 56       | 55          | 49       |
| Tippecanoe             | Kosciusko  | 2018 | 47       | 48          | 43       |

| Lake Name           | County     | Year | TSI [SD] | TSI [Chl-a] | TSI [TP] |
|---------------------|------------|------|----------|-------------|----------|
| Twin Pitts, East    | Pike       | 2018 | 42       | 35          | 55       |
| Twin Pitts, West    | Pike       | 2018 | 49       | 42          | 58       |
| University          | Monroe     | 2018 | 42       | 43          | 65       |
| Upper Long          | Noble      | 2018 | 57       | 50          | 54       |
| Versailles          | Ripley     | 2018 | 73       | 72          | 73       |
| Waveland            | Montgomery | 2018 | 75       | 79          | 60       |
| Webster             | Kosciusko  | 2018 | 56       | 59          | 51       |
| Woods (Big Blue #3) | Rush       | 2018 | 72       | 74          | 68       |